

P^ATENT COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

BOULT WADE TENNANT
Verulam Gardens
70 Gray's Inn Road
London WC1X 8BT
ROYAUME-UNI

Date of mailing (day/month/year) 16 October 2000 (16.10.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference SCB51337/002	
International application No. PCT/EP99/09710	International filing date (day/month/year) 07 December 1999 (07.12.99)

1. The following indications appeared on record concerning:

☐ the applicant ☐ the inventor ☒ the agent ☐ the common representative

Name and Address BOULT WADE TENNANT 27 Furnival Street London, EC4A 1PQ United Kingdom	State of Nationality	State of Residence
	Telephone No. +44(0)20 7430 7500	
	Facsimile No. +44(0)20 7831 1768	
	Teleprinter No.	

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☐ the person ☐ the name ☒ the address ☐ the nationality ☐ the residence

Name and Address BOULT WADE TENNANT Verulam Gardens 70 Gray's Inn Road London WC1X 8BT United Kingdom	State of Nationality	State of Residence
	Telephone No. +44(0)20 7430 7500	
	Facsimile No. +44(0)20 7430 7600	
	Teleprinter No.	

3. Further observations, if necessary:

4. A copy of this notification has been sent to:

<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned
<input type="checkbox"/> the International Searching Authority	<input checked="" type="checkbox"/> the elected Offices concerned
<input checked="" type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer A. Karkachi Telephone No.: (41-22) 338.83.38
--	--

INTERNATIONAL COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C.20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 30 August 2000 (30.08.00)	
International application No. PCT/EP99/09710	Applicant's or agent's file reference SCB51337/002
International filing date (day/month/year) 07 December 1999 (07.12.99)	Priority date (day/month/year) 07 December 1998 (07.12.98)
Applicant KALETTA, Titus et al	

1. The designated Office is hereby notified of its election made:



in the demand filed with the International Preliminary Examining Authority on:

07 July 2000 (07.07.00)



in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer A. Karkachi</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	---

15

ATENT COOPERATION TREATY
PCT

REC'D 08 FEB 2001	
WIPO	PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference SCB/51337/002		FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)
International application No. PCT/EP99/09710	International filing date (day/month/year) 07/12/1999	Priority date (day/month/year) 07/12/1998
International Patent Classification (IPC) or national classification and IPC C12N1/04		
Applicant DEVGEN NV et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 9 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☒ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 07/07/2000	Date of completion of this report 05.02.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Sprinks, M Telephone No. +49 89 2399 8706 

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP99/09710

I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

Description, pages:

1-49 as originally filed

Claims, No.:

1-115 as originally filed

Drawings, sheets:

1/2-2/2 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP99/09710

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

see separate sheet

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

☐ the entire international application.

☒ claims Nos. 26-70,73,74,82-86,90-95,114,115.

because:

☐ the said international application, or the said claims Nos. relate to the following subject matter which does not require an international preliminary examination (*specify*):

☒ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. 82-86,114,115 are so unclear that no meaningful opinion could be formed (*specify*):
see separate sheet

☐ the claims, or said claims Nos. are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for the said claims Nos. 26-70,73,74,90-95.

2. A meaningful international preliminary examination report cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

☐ restricted the claims.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP99/09710

- ☐ paid additional fees.
- ☐ paid additional fees under protest.
- ☒ neither restricted nor paid additional fees.
2. ☐ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.
3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is
- ☐ complied with.
- ☒ not complied with for the following reasons:
see separate sheet
4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:
- ☐ all parts.
- ☒ the parts relating to claims Nos. 1-25,71,72,75-89,96-115.

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	
	No:	Claims	1,5,7-10,19,23-25,71,72,75-79,81,87-89,96-103
Inventive step (IS)	Yes:	Claims	
	No:	Claims	2-4,6,11-18,20-22,80,104-112
Industrial applicability (IA)	Yes:	Claims	1-25,71,72,75-81,87-89,96-112
	No:	Claims	

2. Citations and explanations **see separate sheet**

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/09710

The following documents (D) are mentioned for the first time in this opinion/report; the numbering will be adhered to in the rest of the procedure:

- D1: WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23)
- D2: KATSURA ET AL.: 'Isolation, characterization and epistasis of fluoride-resistant mutants of *Caenorhabditis elegans*' GENETICS, vol. 136, 1994, pages 145-154, XP000886900
- D3: VAN SWINDEREN ET AL.: 'Quantitative trait loci controlling halothane sensitivity in *Caenorhabditis elegans*' PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784
- D4: AHRINGER ET AL.: 'Turn to the worm!' CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application

I) Basis of the opinion/report

Additional observations

- 1) The applicant has waived his right to a written opinion and requested an immediate international preliminary examination report.

III) Non-establishment of opinion

Clarity

- 1) Because the subject-matter of **claims 82-86, 114 and 115** is so unclear, a meaningful assessment of novelty/inventive step cannot be made at the present time. However, in order to expedite the procedure, the applicant is requested to note the following points:
- 2) Although said claims are directed to methods for elucidating biochemical pathways in a nematode worm, the defining steps of said methods appear merely to result in gross phenotypic comparisons between different genetic defects (in any event, the subject-matter of said claims would not be considered novel or

inventive for similar reasons to those given in section V below).

- 3) **Claims 114 and 115** refer broadly to libraries of nematode worms which, in the absence of any other technical features would be indistinguishable from collections of nematode worms in general (and, therefore, also not new).

IV) Unity

- 1) This authority made an objection concerning lack of unity of invention for the originally filed application which was in agreement with an objection previously put forward by the International Searching Authority (**Rule 13.1 - 13.3 PCT**). The objection is summarised below:
- 2) The following 4 inventions identified within originally filed **claims** are not so linked as to form a single general inventive concept:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81,96-115 partially:

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81,96-115 partially:

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81,96-115 partially: Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially: Method for

determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

- 3) The only common concept linking the above subjects is that of providing libraries of nematodes scored for multiple phenotypic traits for determining the modes of action of different compounds (including genes and their products). However, since other such libraries and uses thereof are disclosed in WO 90 09096 A (see page 7, line 18 - page 8, line 23 and page 15, lines 14-30), this concept is not novel. Consequently, each of the subjects defined above constitutes a separate invention.
- 4) In response to an invitation to restrict the claims or pay additional taxes, the applicant paid no additional taxes but elected invention 1 for substantive examination, to which **claims 1-25, 71, 72, 75-89 and 96-115** correspond.

V) Reasoned statement

Novelty

- 1) The present application does not satisfy the criterion set forth in **Article 33 (2) PCT** because the subject-matter of **claims 1, 5, 7-10, 19, 23-25, 71, 72, 75-79, 81, 87-89 and 96-103** is not new in respect of prior art as defined in the regulations (**Rule 64.1 - 64.3 PCT**).
- 2) D1 discloses methods for screening and classifying compounds of pharmaceutical interest comprising evaluating the phenotypic effect of a compound on a series of *C. elegans* nematodes selected from the group consisting of wild-type, stable mutants or both and comparing said effect with a phenotypic library compiled from the multiple phenotypic effects (e.g. paralysis, egg laying) resulting from exposing said series of nematodes to other (e.g. known) compounds of the prior art (see abstract, pages 7-8 "summary of the invention" and page 15, lines 14-30. The outcome of such methods is the functional/biochemical characterisation of compounds with respect to compounds and/or genes with a known activity (over 700 genetic mutations in *C. elegans* were known at the priority date of D1 - practically the whole genome had been sequenced at the priority date of the

present application).

Consequently, **claims 1, 5, 8, 19, 23-25, 71, 72, 75-79, 81, 87-89 and 96-103** lack novelty in the light of D1.

- 3) D2 discloses the generation of a library of 13 fluoride-resistant *C. elegans* mutants (defining five new genes), phenotypically scored for growth rates and brood sizes. It also discloses the construction of double and triple mutants and their phenotypic comparison with said library to ascertain the genetic/biochemical nature of the different fluoride-resistance mutations (see abstract and especially tables 1-4).

Consequently, **claims 1, 5, 7-10, 19, 23-25, 87-89, 102 and 103** lack novelty in the light of D2 (D3 is similarly relevant).

Inventive Step

- 4) The present application does not satisfy the criterion set forth in **Article 33 (3) PCT** because the subject-matter of **claims 2-4, 6, 11-18, 20-22, 80 and 104-112** does not involve an inventive step (**Rule 65.1 and 65.2 PCT**).
- 5) Each of D1-D3 (especially D1) discloses the general concept of providing *C. elegans* libraries scored for multiple phenotypic traits and their use for the elucidation of compound/gene activities. In the light of this concept, none of the features of said claims, in combination with the features of the claims to which they refer, could be considered to involve an inventive step, since they merely represent obvious alternatives of which a person skilled in the art would be aware (see D4 for a review article highlighting such obvious alternatives).

VIII) Certain observations

Clarity

- 1) The present application does not satisfy the criterion set forth in **Article 6 PCT** because the subject-matter of the claims in general is unclear.

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/09710

- 2) Many of the methods claimed are formulated in such a broad manner that their subject-matters substantially overlap, making it extremely difficult to determine the essential technical features in each case.
- 3) Since even a single or a few worms may be considered to be a library, many of the claims (see **claim 71** for example) are also unclear because they may be considered to encompass screening compounds against a worm with a genetic defect in order to find a compound capable of restoring the wild-type phenotype (comparison with a "wild-type phenotypic library"). Such an interpretation could lead to further novelty/inventive step objections at a later date.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT
(PCT Article 36 and Rule 70)



Applicant's or agent's file reference SCB/51337/002	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP99/09710	International filing date (day/month/year) 07/12/1999	Priority date (day/month/year) 07/12/1998
International Patent Classification (IPC) or national classification and IPC C12N1/04		
Applicant DEVGEN NV et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 9 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:
 - I ☒ Basis of the report
 - II ☐ Priority
 - III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
 - IV ☒ Lack of unity of invention
 - V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
 - VI ☐ Certain documents cited
 - VII ☐ Certain defects in the international application
 - VIII ☒ Certain observations on the international application

Date of submission of the demand 07/07/2000	Date of completion of this report 05.02.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Sprinks, M Telephone No. +49 89 2399 8706 

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP99/09710

I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

Description, pages:

1-49 as originally filed

Claims, No.:

1-115 as originally filed

Drawings, sheets:

1/2-2/2 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP99/09710

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:
see separate sheet

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

☐ the entire international application.

☒ claims Nos. 26-70,73,74,82-86,90-95,114,115.

because:

☐ the said international application, or the said claims Nos. relate to the following subject matter which does not require an international preliminary examination (*specify*):

☒ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. 82-86,114,115 are so unclear that no meaningful opinion could be formed (*specify*):
see separate sheet

☐ the claims, or said claims Nos. are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for the said claims Nos. 26-70,73,74,90-95.

2. A meaningful international preliminary examination report cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

☐ restricted the claims.

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP99/09710

- ☐ paid additional fees.
- ☐ paid additional fees under protest.
- ☒ neither restricted nor paid additional fees.
2. ☐ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.
3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is
- ☐ complied with.
- ☒ not complied with for the following reasons:
see separate sheet
4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:
- ☐ all parts.
- ☒ the parts relating to claims Nos. 1-25,71,72,75-89,96-115.

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims
	No:	Claims 1,5,7-10,19,23-25,71,72,75-79,81,87-89,96-103
Inventive step (IS)	Yes:	Claims
	No:	Claims 2-4,6,11-18,20-22,80,104-112
Industrial applicability (IA)	Yes:	Claims 1-25,71,72,75-81,87-89,96-112
	No:	Claims

2. Citations and explanations
see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/09710

The following documents (D) are mentioned for the first time in this opinion/report; the numbering will be adhered to in the rest of the procedure:

- D1: WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23)
- D2: KATSURA ET AL.: 'Isolation, characterization and epistasis of fluoride-resistant mutants of *Caenorhabditis elegans*' GENETICS, vol. 136, 1994, pages 145-154, XP000886900
- D3: VAN SWINDEREN ET AL.: 'Quantitative trait loci controlling halothane sensitivity in *Caenorhabditis elegans*' PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784
- D4: AHRINGER ET AL.: 'Turn to the worm!' CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application

I) Basis of the opinion/report

Additional observations

- 1) The applicant has waived his right to a written opinion and requested an immediate international preliminary examination report.

III) Non-establishment of opinion

Clarity

- 1) Because the subject-matter of **claims 82-86, 114 and 115** is so unclear, a meaningful assessment of novelty/inventive step cannot be made at the present time. However, in order to expedite the procedure, the applicant is requested to note the following points:
- 2) Although said claims are directed to methods for elucidating biochemical pathways in a nematode worm, the defining steps of said methods appear merely to result in gross phenotypic comparisons between different genetic defects (in any event, the subject-matter of said claims would not be considered novel or

inventive for similar reasons to those given in section V below).

- 3) **Claims 114 and 115** refer broadly to libraries of nematode worms which, in the absence of any other technical features would be indistinguishable from collections of nematode worms in general (and, therefore, also not new).

IV) Unity

- 1) This authority made an objection concerning lack of unity of invention for the originally filed application which was in agreement with an objection previously put forward by the International Searching Authority (**Rule 13.1 - 13.3 PCT**). The objection is summarised below:
- 2) The following 4 inventions identified within originally filed **claims** are not so linked as to form a single general inventive concept:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81,96-115 partially:

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81,96-115 partially:

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81,96-115 partially: Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially: Method for

determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

- 3) The only common concept linking the above subjects is that of providing libraries of nematodes scored for multiple phenotypic traits for determining the modes of action of different compounds (including genes and their products). However, since other such libraries and uses thereof are disclosed in WO 90 09096 A (see page 7, line 18 - page 8, line 23 and page 15, lines 14-30), this concept is not novel. Consequently, each of the subjects defined above constitutes a separate invention.
- 4) In response to an invitation to restrict the claims or pay additional taxes, the applicant paid no additional taxes but elected invention 1 for substantive examination, to which **claims 1-25, 71, 72, 75-89 and 96-115** correspond.

V) Reasoned statement

Novelty

- 1) The present application does not satisfy the criterion set forth in **Article 33 (2) PCT** because the subject-matter of **claims 1, 5, 7-10, 19, 23-25, 71, 72, 75-79, 81, 87-89 and 96-103** is not new in respect of prior art as defined in the regulations (**Rule 64.1 - 64.3 PCT**).
- 2) D1 discloses methods for screening and classifying compounds of pharmaceutical interest comprising evaluating the phenotypic effect of a compound on a series of *C. elegans* nematodes selected from the group consisting of wild-type, stable mutants or both and comparing said effect with a phenotypic library compiled from the multiple phenotypic effects (e.g. paralysis, egg laying) resulting from exposing said series of nematodes to other (e.g. known) compounds of the prior art (see abstract, pages 7-8 "summary of the invention" and page 15, lines 14-30. The outcome of such methods is the functional/biochemical characterisation of compounds with respect to compounds and/or genes with a known activity (over 700 genetic mutations in *C. elegans* were known at the priority date of D1 - practically the whole genome had been sequenced at the priority date of the

present application).

Consequently, **claims 1, 5, 8, 19, 23-25, 71, 72, 75-79, 81, 87-89 and 96-103** lack novelty in the light of D1.

- 3) D2 discloses the generation of a library of 13 fluoride-resistant *C. elegans* mutants (defining five new genes), phenotypically scored for growth rates and brood sizes. It also discloses the construction of double and triple mutants and their phenotypic comparison with said library to ascertain the genetic/biochemical nature of the different fluoride-resistance mutations (see abstract and especially tables 1-4).

Consequently, **claims 1, 5, 7-10, 19, 23-25, 87-89, 102 and 103** lack novelty in the light of D2 (D3 is similarly relevant).

Inventive Step

- 4) The present application does not satisfy the criterion set forth in **Article 33 (3) PCT** because the subject-matter of **claims 2-4, 6, 11-18, 20-22, 80 and 104-112** does not involve an inventive step (**Rule 65.1 and 65.2 PCT**).
- 5) Each of D1-D3 (especially D1) discloses the general concept of providing *C. elegans* libraries scored for multiple phenotypic traits and their use for the elucidation of compound/gene activities. In the light of this concept, none of the features of said claims, in combination with the features of the claims to which they refer, could be considered to involve an inventive step, since they merely represent obvious alternatives of which a person skilled in the art would be aware (see D4 for a review article highlighting such obvious alternatives).

VIII) Certain observations

Clarity

- 1) The present application does not satisfy the criterion set forth in **Article 6 PCT** because the subject-matter of the claims in general is unclear.

- 2) Many of the methods claimed are formulated in such a broad manner that their subject-matters substantially overlap, making it extremely difficult to determine the essential technical features in each case.
- 3) Since even a single or a few worms may be considered to be a library, many of the claims (see **claim 71** for example) are also unclear because they may be considered to encompass screening compounds against a worm with a genetic defect in order to find a compound capable of restoring the wild-type phenotype (comparison with a "wild-type phenotypic library"). Such an interpretation could lead to further novelty/inventive step objections at a later date.

PATENT COOPERATION TREATY

From the INTERNATIONAL SEARCHING AUTHORITY

PCT

To:
BOULT WADE TENNANT
27 Furnival Street
London EC4A 1PQ
UNITED KINGDOM

NOTIFICATION OF TRANSMITTAL OF
THE INTERNATIONAL SEARCH REPORT
OR THE DECLARATION

(PCT Rule 44.1)

RECEIVED

31 JUL 2000

<p>Applicant's or agent's file reference SCB51337/002</p> <p>International application No. PCT/EP 99/ 09710</p> <p>Applicant DEVGEN NV et al.</p>	<p>Date of mailing (day/month/year) 25/07/2000</p> <p>FOR FURTHER ACTION See paragraphs 1 and 4 below</p> <p>International filing date (day/month/year) 07/12/1999</p>
--	---

1. ☒ The applicant is hereby notified that the International Search Report has been established and is transmitted herewith.

Filing of amendments and statement under Article 19:
The applicant is entitled, if he so wishes, to amend the claims of the International Application (see Rule 46):

When? The time limit for filing such amendments is normally 2 months from the date of transmittal of the International Search Report; however, for more details, see the notes on the accompanying sheet.

Where? Directly to the International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland
Facsimile No.: (41-22) 740.14.35

For more detailed instructions, see the notes on the accompanying sheet.

2. ☐ The applicant is hereby notified that no International Search Report will be established and that the declaration under Article 17(2)(a) to that effect is transmitted herewith.

3. ☐ **With regard to the protest** against payment of (an) additional fee(s) under Rule 40.2, the applicant is notified that:

☐ the protest together with the decision thereon has been transmitted to the International Bureau together with the applicant's request to forward the texts of both the protest and the decision thereon to the designated Offices.

☐ no decision has been made yet on the protest; the applicant will be notified as soon as a decision is made.

4. **Further action(s):** The applicant is reminded of the following:

Shortly after **18 months** from the priority date, the international application will be published by the International Bureau. If the applicant wishes to avoid or postpone publication, a notice of withdrawal of the international application, or of the priority claim, must reach the International Bureau as provided in Rules 90bis.1 and 90bis.3, respectively, before the completion of the technical preparations for international publication.

Within **19 months** from the priority date, a demand for international preliminary examination must be filed if the applicant wishes to postpone the entry into the national phase until 30 months from the priority date (in some Offices even later).

Within **20 months** from the priority date, the applicant must perform the prescribed acts for entry into the national phase before all designated Offices which have not been elected in the demand or in a later election within 19 months from the priority date or could not be elected because they are not bound by Chapter II.

<p>Name and mailing address of the International Searching Authority</p> <p> European Patent Office, P.B. 5818 Patentlaan 2 NL-2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016</p>	<p>Authorized officer</p> <p>Véronique Bafflou</p>
---	---

NOTES TO FORM PCT/ISA/220

These Notes are intended to give the basic instructions concerning the filing of amendments under article 19. The Notes are based on the requirements of the Patent Cooperation Treaty, the Regulations and the Administrative Instructions under that Treaty. In case of discrepancy between these Notes and those requirements, the latter are applicable. For more detailed information, see also the PCT Applicant's Guide, a publication of WIPO.

In these Notes, "Article", "Rule", and "Section" refer to the provisions of the PCT, the PCT Regulations and the PCT Administrative Instructions, respectively.

INSTRUCTIONS CONCERNING AMENDMENTS UNDER ARTICLE 19

The applicant has, after having received the international search report, one opportunity to amend the claims of the international application. It should however be emphasized that, since all parts of the international application (claims, description and drawings) may be amended during the international preliminary examination procedure, there is usually no need to file amendments of the claims under Article 19 except where, e.g. the applicant wants the latter to be published for the purposes of provisional protection or has another reason for amending the claims before international publication. Furthermore, it should be emphasized that provisional protection is available in some States only.

What parts of the international application may be amended?

Under Article 19, only the claims may be amended.

During the international phase, the claims may also be amended (or further amended) under Article 34 before the International Preliminary Examining Authority. The description and drawings may only be amended under Article 34 before the International Examining Authority.

Upon entry into the national phase, all parts of the international application may be amended under Article 28 or, where applicable, Article 41.

When?

Within 2 months from the date of transmittal of the international search report or 16 months from the priority date, whichever time limit expires later. It should be noted, however, that the amendments will be considered as having been received on time if they are received by the International Bureau after the expiration of the applicable time limit but before the completion of the technical preparations for international publication (Rule 46.1).

Where not to file the amendments?

The amendments may only be filed with the International Bureau and not with the receiving Office or the International Searching Authority (Rule 46.2).

Where a demand for international preliminary examination has been/is filed, see below.

How?

Either by cancelling one or more entire claims, by adding one or more new claims or by amending the text of one or more of the claims as filed.

A replacement sheet must be submitted for each sheet of the claims which, on account of an amendment or amendments, differs from the sheet originally filed.

All the claims appearing on a replacement sheet must be numbered in Arabic numerals. Where a claim is cancelled, no renumbering of the other claims is required. In all cases where claims are renumbered, they must be renumbered consecutively (Administrative Instructions, Section 205(b)).

The amendments must be made in the language in which the international application is to be published.

What documents must/may accompany the amendments?

Letter (Section 205(b)):

The amendments must be submitted with a letter.

The letter will not be published with the international application and the amended claims. It should not be confused with the "Statement under Article 19(1)" (see below, under "Statement under Article 19(1)").

The letter must be in English or French, at the choice of the applicant. However, if the language of the international application is English, the letter must be in English; if the language of the international application is French, the letter must be in French.

NOTES TO FORM PCT/ISA/220 (continued)

The letter must indicate the differences between the claims as filed and the claims as amended. It must, in particular, indicate, in connection with each claim appearing in the international application (it being understood that identical indications concerning several claims may be grouped), whether

- (i) the claim is unchanged;
- (ii) the claim is cancelled;
- (iii) the claim is new;
- (iv) the claim replaces one or more claims as filed;
- (v) the claim is the result of the division of a claim as filed.

The following examples illustrate the manner in which amendments must be explained in the accompanying letter:

1. [Where originally there were 48 claims and after amendment of some claims there are 51]:
"Claims 1 to 29, 31, 32, 34, 35, 37 to 48 replaced by amended claims bearing the same numbers; claims 30, 33 and 36 unchanged; new claims 49 to 51 added."
2. [Where originally there were 15 claims and after amendment of all claims there are 11]:
"Claims 1 to 15 replaced by amended claims 1 to 11."
3. [Where originally there were 14 claims and the amendments consist in cancelling some claims and in adding new claims]:
"Claims 1 to 6 and 14 unchanged; claims 7 to 13 cancelled; new claims 15, 16 and 17 added." or
"Claims 7 to 13 cancelled; new claims 15, 16 and 17 added; all other claims unchanged."
4. [Where various kinds of amendments are made]:
"Claims 1-10 unchanged; claims 11 to 13, 18 and 19 cancelled; claims 14, 15 and 16 replaced by amended claim 14; claim 17 subdivided into amended claims 15, 16 and 17; new claims 20 and 21 added."

"Statement under article 19(1)" (Rule 46.4)

The amendments may be accompanied by a statement explaining the amendments and indicating any impact that such amendments might have on the description and the drawings (which cannot be amended under Article 19(1)).

The statement will be published with the international application and the amended claims.

It must be in the language in which the international application is to be published.

It must be brief, not exceeding 500 words if in English or if translated into English.

It should not be confused with and does not replace the letter indicating the differences between the claims as filed and as amended. It must be filed on a separate sheet and must be identified as such by a heading, preferably by using the words "Statement under Article 19(1)."

It may not contain any disparaging comments on the international search report or the relevance of citations contained in that report. Reference to citations, relevant to a given claim, contained in the international search report may be made only in connection with an amendment of that claim.

Consequence if a demand for international preliminary examination has already been filed

If, at the time of filing any amendments and any accompanying statement, under Article 19, a demand for international preliminary examination has already been submitted, the applicant must preferably, at the time of filing the amendments (and any statement) with the International Bureau, also file with the International Preliminary Examining Authority a copy of such amendments (and of any statement) and, where required, a translation of such amendments for the procedure before that Authority (see Rules 55.3(a) and 62.2, first sentence). For further information, see the Notes to the demand form (PCT/IPEA/401).

Consequence with regard to translation of the international application for entry into the national phase

The applicant's attention is drawn to the fact that, upon entry into the national phase, a translation of the claims as amended under Article 19 may have to be furnished to the designated/elected Offices, instead of, or in addition to, the translation of the claims as filed.

For further details on the requirements of each designated/elected Office, see Volume II of the PCT Applicant's Guide.

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference SCB51337/002	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 99/ 09710	International filing date (day/month/year) 07/12/1999	(Earliest) Priority Date (day/month/year) 07/12/1998
Applicant DEVGEN NV et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 7 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☒ **Unity of invention is lacking** (see Box II).

4. With regard to the title,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the abstract,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 99/09710

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 113
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 113

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially

Method for determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09710

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N1/04 C12N1/00 C12N15/01 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23) Cited against inventions 1 and 2 in their entirety and inventions 3 and 4 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. page 7, line 18 -page 8, line 23 page 15, line 14 - line 30 --- -/--</p>	<p>1-112, 114,115</p>

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

25.07.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KATSURA ET AL.: "Isolation, characterization and epistasis of fluoride-resistant mutants of <i>Caenorhabditis elegans</i> " GENETICS, vol. 136, 1994, pages 145-154, XP000886900 Cited against invention 1 abstract; tables 1-4 page 145, column 1 -page 146, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
X	VAN SWINDEREN ET AL.: "Quantitative trait loci controlling halothane sensitivity in <i>Caenorhabditis elegans</i> " PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784 Cited against invention 2 in its entirety and invention 3 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. abstract page 8232, column 1 -page 8233, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
A	AHRINGER ET AL.: "Turn to the worm!" CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application Cited for all inventions the whole document ---	1-112, 114,115
X	WO 96 38555 A (BOGAERT THIERRY ;STRINGHAM EVE (CA); VANDEKERCKHOVE JOEL (BE)) 5 December 1996 (1996-12-05) Cited against inventions 2 and 3 page 35, line 22 -page 36, line 28; claim 43 ---	26-68, 71-77, 79-81, 90-114
A	SAMOILOFF, M.R. ET AL: "The use of nematodes in marine ecotoxicology. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 1." MAR. TOX., (1984) PP. 407-426. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000812-8., XP000886947 Dep. Zool., Univ. Manitoba, Winnipeg, Man. R3T 2N2, Canada Cited for inventions 3 and 4 page 413, paragraph 2 --- -/--	55-68, 71,72, 75, 79-81, 93-112, 114,115

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BOGAERT, T. ET AL: "Determination of the toxicity of four heavy metal compounds and three carcinogens using two marine nematode species, <i>Monhystera microphthalma</i> and <i>Diploelaimelloides brucei</i>. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 2." MAR. TOX., (1984) PP. 21-30. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4;;90-9000813-6., XP000886948</p> <p>Lab. Mol. Biol., Med. Res. Counc. Cent., University Med. Sch., Hills Rd., Cambridge CB2 2QH, UK</p> <p>Cited for inventions 3 and 4 the whole document</p> <p>-----</p>	<p>55-68, 71,72, 75, 79-81, 93-112, 114,115</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/09710

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9009096	A	23-08-1990	AU	5106790 A	05-09-1990
WO 9638555	A	05-12-1996	AU	6123496 A	18-12-1996
			EP	0832222 A	01-04-1998

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference SCB51337/002	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 99/09710	International filing date (day/month/year) 07/12/1999	(Earliest) Priority Date (day/month/year) 07/12/1998
Applicant DEVGEN NV et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 7 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☒ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 99/09710

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 113
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 113

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially

Method for determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N1/04 C12N15/00 C12N15/01 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23) Cited against inventions 1 and 2 in their entirety and inventions 3 and 4 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. page 7, line 18 -page 8, line 23 page 15, line 14 - line 30 --- -/--</p>	<p>1-112, 114,115</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

25. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	KATSURA ET AL.: "Isolation, characterization and epistasis of fluoride-resistant mutants of <i>Caenorhabditis elegans</i> " GENETICS, vol. 136, 1994, pages 145-154, XP000886900 Cited against invention 1 abstract; tables 1-4 page 145, column 1 -page 146, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
X	VAN SWINDEREN ET AL.: "Quantitative trait loci controlling halothane sensitivity in <i>Caenorhabditis elegans</i> " PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784 Cited against invention 2 in its entirety and invention 3 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. abstract page 8232, column 1 -page 8233, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
A	AHRINGER ET AL.: "Turn to the worm!" CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application Cited for all inventions the whole document ---	1-112, 114,115
X	WO 96 38555 A (BOGAERT THIERRY ;STRINGHAM EVE (CA); VANDEKERCKHOVE JOEL (BE)) 5 December 1996 (1996-12-05) Cited against inventions 2 and 3 page 35, line 22 -page 36, line 28; claim 43 ---	26-68, 71-77, 79-81, 90-114
A	SAMOILOFF, M.R. ET AL: "The use of nematodes in marine ecotoxicology. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 1." MAR. TOX., (1984) PP. 407-426. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4;;90-9000812-8., XP000886947 Dep. Zool., Univ. Manitoba, Winnipeg, Man. R3T 2N2, Canada Cited for inventions 3 and 4 page 413, paragraph 2 ---	55-68, 71,72, 75, 79-81, 93-112, 114,115

-/--

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BOGAERT, T. ET AL: "Determination of the toxicity of four heavy metal compounds and three carcinogens using two marine nematode species, Monhystera microphthalma and Diplolaimelloides brucei. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 2." MAR. TOX., (1984) PP. 21-30. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000813-6., XP000886948</p> <p>Lab. Mol. Biol., Med. Res. Counc. Cent., University Med. Sch., Hills Rd., Cambridge CB2 2QH, UK</p> <p>Cited for inventions 3 and 4 the whole document</p> <p>-----</p>	<p>55-68, 71,72, 75, 79-81, 93-112, 114,115</p>

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9009096	A	23-08-1990	AU	5106790 A	05-09-1990

WO 9638555	A	05-12-1996	AU	6123496 A	18-12-1996
			EP	0832222 A	01-04-1998

2L
PCT

INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 1/04, 1/00, 15/01, 15/10		A3	(11) International Publication Number: WO 00/34438
			(43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/EP99/09710 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 9826890.7 7 December 1998 (07.12.98) GB (71) Applicant (for all designated States except US): DEVGEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KALETTA, Titus [BE/BE]; (BE). FEICHTINGER, Richard [BE/BE]; (BE). VAN POUCKE, Jonas [BE/BE]; (BE). VAN GEEL, Anton [BE/BE]; (BE). APPELMANS, Saskia [BE/BE]; (BE). VAN CRIEKINGE, Wim [BE/BE]; (BE). BOGAERT, Thierry [BE/BE]; Devgen NV, Technologiepark 9, B-9052 Zwijnaarde (BE). (74) Agent: BOULT WADE TENNANT; 27 Fumival Street, London, EC4A 1PQ (GB).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 9 November 2000 (09.11.00)
(54) Title: METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC PROFILES			
(57) Abstract Methods are provided for use in constructing libraries of phenotypic profiles in a nematode worm such as <i>C. elegans</i> . The methods require measurement of identifiable characteristics of the worm and systematic scoring of these characteristics. Also provided are methods of identifying compounds with potential pharmacological activity, for determining the mode of action of a given compound and for assigning genes to particular biochemical pathways.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/99/09710

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N1/04 C12N1/00 C12N15/01 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

X	<p>WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23) Cited against inventions 1 and 2 in their entirety and inventions 3 and 4 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. page 7, line 18 -page 8, line 23 page 15, line 14 - line 30 --- -/--</p>	<p>1-112, 114,115</p>
---	---	---------------------------

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

25. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/E/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KATSURA ET AL.: "Isolation, characterization and epistasis of fluoride-resistant mutants of <i>Caenorhabditis elegans</i> " GENETICS, vol. 136, 1994, pages 145-154, XP000886900 Cited against invention 1 abstract; tables 1-4 page 145, column 1 -page 146, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
X	VAN SWINDEREN ET AL.: "Quantitative trait loci controlling halothane sensitivity in <i>Caenorhabditis elegans</i> " PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784 Cited against invention 2 in its entirety and invention 3 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. abstract page 8232, column 1 -page 8233, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
A	AHRINGER ET AL.: "Turn to the worm!" CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application Cited for all inventions the whole document ---	1-112, 114,115
X	WO 96 38555 A (BOGAERT THIERRY ;STRINGHAM EVE (CA); VANDEKERCKHOVE JOEL (BE)) 5 December 1996 (1996-12-05) Cited against inventions 2 and 3 page 35, line 22 -page 36, line 28; claim 43 ---	26-68, 71-77, 79-81, 90-114
A	SAMOILOFF, M.R. ET AL: "The use of nematodes in marine ecotoxicology. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 1." MAR. TOX., (1984) PP. 407-426. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000812-8., XP000886947 Dep. Zool., Univ. Manitoba, Winnipeg, Man. R3T 2N2, Canada Cited for inventions 3 and 4 page 413, paragraph 2 ---	55-68, 71,72, 75, 79-81, 93-112, 114,115

-/--

INTERNATIONAL SEARCH REPORT

international application no

PCT/E 9/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BOGAERT, T. ET AL: "Determination of the toxicity of four heavy metal compounds and three carcinogens using two marine nematode species, Monhystera microphthalma and Diplolaimelloides brucei. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 2."</p> <p>MAR. TOX., (1984) PP. 21-30. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4;;90-9000813-6., XP000886948</p> <p>Lab. Mol. Biol., Med. Res. Counc. Cent., University Med. Sch., Hills Rd., Cambridge CB2 2QH, UK</p> <p>Cited for inventions 3 and 4 the whole document</p> <p>-----</p>	<p>55-68, 71,72, 75, 79-81, 93-112, 114,115</p>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 99/09710

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 113
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially

Method for determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 113

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/EP 99/09710

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
W0 9009096	A	23-08-1990	AU	5106790 A	05-09-1990

W0 9638555	A	05-12-1996	AU	6123496 A	18-12-1996
			EP	0832222 A	01-04-1998

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 1/04, 1/00, 15/01, 15/10	A3	(11) International Publication Number: WO 00/34438 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/EP99/09710 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 9826890.7 7 December 1998 (07.12.98) GB (71) Applicant (for all designated States except US): DEVGEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KALETTA, Titus [BE/BE]; (BE). FEICHTINGER, Richard [BE/BE]; (BE). VAN POUCKE, Jonas [BE/BE]; (BE). VAN GEEL, Anton [BE/BE]; (BE). APPELMANS, Saskia [BE/BE]; (BE). VAN CRIEKINGE, Wim [BE/BE]; (BE). BOGAERT, Thierry [BE/BE]; Devgen NV, Technologiepark 9, B-9052 Zwijnaarde (BE). (74) Agent: BOULT WADE TENNANT; 27 Fumival Street, London, EC4A 1PQ (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 9 November 2000 (09.11.00)
(54) Title: METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC PROFILES		
(57) Abstract Methods are provided for use in constructing libraries of phenotypic profiles in a nematode worm such as <i>C. elegans</i> . The methods require measurement of identifiable characteristics of the worm and systematic scoring of these characteristics. Also provided are methods of identifying compounds with potential pharmacological activity, for determining the mode of action of a given compound and for assigning genes to particular biochemical pathways.		

INTERNATIONAL SEARCH REPORT

International Application No.

EP 99/09710

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N1/04 C12N1/00 C12N15/01 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 90 09096 A. (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23) Cited against inventions 1 and 2 in their entirety and inventions 3 and 4 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. page 7, line 18 -page 8, line 23 page 15, line 14 - line 30 --- -/--</p>	1-112, 114,115

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

25. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

INTERNATIONAL SEARCH REPORT

International Application No.
P 99/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BOGAERT, T. ET AL: "Determination of the toxicity of four heavy metal compounds and three carcinogens using two marine nematode species, <i>Monhystera microphthalma</i> and <i>Diploelaimeloides brucei</i>. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 2." MAR. TOX., (1984) PP. 21-30. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000813-6., XP000886948</p> <p>Lab. Mol. Biol., Med. Res. Counc. Cent., University Med. Sch., Hills Rd., Cambridge CB2 2QH, UK</p> <p>Cited for inventions 3 and 4 the whole document</p> <p>-----</p>	<p>55-68, 71,72, 75, 79-81, 93-112, 114,115</p>

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially

Method for determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/EP 99/09710

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9009096	A	23-08-1990	AU	5106790 A	05-09-1990

WO 9638555	A	05-12-1996	AU	6123496 A	18-12-1996
			EP	0832222 A	01-04-1998



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 1/04, 1/00, 15/01, 15/10	A2	(11) International Publication Number: WO 00/34438 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/EP99/09710 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 9826890.7 7 December 1998 (07.12.98) GB (71) Applicant (for all designated States except US): DEVGEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KALETTA, Titus [BE/BE]; (BE). FEICHTINGER, Richard [BE/BE]; (BE). VAN POUCKE, Jonas [BE/BE]; (BE). VAN GEEL, Anton [BE/BE]; (BE). APPELMANS, Saskia [BE/BE]; (BE). VAN CRIEKINGE, Wim [BE/BE]; (BE). BOGAERT, Thierry [BE/BE]; Devgen NV, Technologiepark 9, B-9052 Zwijnaarde (BE). (74) Agent: BOULT WADE TENNANT; 27 Furnival Street, London, EC4A 1PQ (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC PROFILES		
(57) Abstract <p>Methods are provided for use in constructing libraries of phenotypic profiles in a nematode worm such as <i>C. elegans</i>. The methods require measurement of identifiable characteristics of the worm and systematic scoring of these characteristics. Also provided are methods of identifying compounds with potential pharmacological activity, for determining the mode of action of a given compound and for assigning genes to particular biochemical pathways.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC
PROFILES

The present invention is concerned with the field
5 of 'genetic pharmacology'. Specifically, it relates to
methods which can determine, among other things,
whether a compound has potential pharmacological
activity, whether a compound interacts with a
particular gene or biochemical pathway in man or
10 animals, what side effects are likely to be associated
with a particular pharmaceutical compound and/or the
mode or modes of action of any compound with
biological activity. Additional uses for the methods
of the invention include the assignment of function to
15 particular genes or assignment of genes and their
encoded proteins to particular biochemical pathways.
In particular, the invention relates to the use of a
microscopic nematode worm, for example *Caenorhabditis
elegans*, and libraries of such worms in the
20 aforementioned methods. These new methods are able to
enhance and accelerate the drug discovery process.

Prior to the early 1990's the search for new
compounds having the potential to combat human or
animal disease was often begun by taking a compound
25 known to have a particular pharmacological activity,
synthesising structurally related variants and then
testing those variants against the known target.

The test against the target might be carried out
in vivo, for example by use of animal models of a
30 human disease. Alternatively, if a particular molecule
was known to be implicated in the progress of a
disease, the compounds could be tested for
interaction with the molecule *in vitro*. The
limitations of such methods are that in the event of a
35 negative result no other information about the
pharmaceutical potential of the compound tested is

- 2 -

gained. For example, an *in vitro* test might show a compound to have no inhibitory action against a particular target enzyme but that compound might have an inhibitory action against another enzyme in the same biochemical pathway as the target enzyme and therefore, in fact, have potential in treatment of the target disease. Animal tests, while providing a reasonable indication of both efficacy and toxicity, provide no information at all about the mode of action of the compound, and therefore the possible reasons for any toxicity. Furthermore, they are time-consuming and expensive and do not lend themselves to automation.

Since the early nineties there have been two developments in particular which have revolutionized the drug discovery process, these being the new sciences of 'genomics' and 'combinatorial chemistry'. It has now been realised that a vast number of diseases have a genetic component and they are not purely the result of environmental influences. Indeed, it is possible that nearly all diseases are multifactorial and will have some degree of genetic basis, albeit very small in some cases. A huge amount of effort is being directed at the present time to the study of the organisation of the genomes of various unicellular and multicellular organisms, including humans. This involves the identification and sequencing of all the genes in a particular genome. Such activity does not only allow for hunting of genes which are directly associated with particular diseases but each of the genes found and the proteins they encode can become, directly or indirectly, a target against which compounds can be screened, whether or not that gene has yet been associated with a disease or indeed has any identified function at all.

- 3 -

Furthermore, rather than starting from a compound of known 'activity' and relying on theoretical structure/function relationships to synthesise new candidate compounds, vast libraries of compounds, of uniform activity can be very rapidly synthesized in an automated manner by combinatorial chemistry. Thus, there is now potential to screen thousands of compounds against thousands of genes and the proteins they encode in very rapid high throughput screens (HTS) and to link compounds to genes and genes to disease.

The present inventors have discovered that these new technologies for drug discovery can conveniently be married with a particular multicellular organism, a nematode worm, *C.elegans*, which has been well characterised genetically and morphologically. They have thereby developed new methods, which are extremely powerful, rapid and convenient and can play an essential part in a drug discovery program.

C. elegans is a microscopic nematode worm which occurs naturally in the soil but can be easily grown in the laboratory on nutrient agar inoculated with bacteria, preferably *E. coli*, on which it feeds. Each worm grows from an embryo to an adult worm of about 1 mm long in three days or so. As it is fully transparent at all stages of its life, cell divisions, migrations and differentiation can be seen in live animals. Furthermore, although its anatomy is simple its somatic cells represent most major differentiated tissue type including muscles, neurons, intestine and epidermis. Accordingly, differences in phenotype which represent a departure from that of a wild-type worm are relatively easily observed, either directly by microscopy or by using selective staining procedures, and many of these phenotypic differences submit to quantitative measurement. Many *C. elegans* mutants have

- 4 -

been identified and their phenotypes described, for example, see *C. elegans* II Ed. Riddle, Blumenthal, Meyer and Priess, Cold Spring Harbor Laboratory Press, 1997. The *C. elegans* genome is now almost entirely
5 sequenced as a result of the *C. elegans* genome project, carried out at the Sanger Center and Washington University School of Medicine. The sequence is available in a public database at
10 http://www.sanger.ac.uk/projects/C_elegans/. As a result of this it has emerged that *C. elegans* comprises genes which have equivalents that are widely distributed in most or all animals including humans.

Methods for creating mutant worms with mutations in selected *C. elegans* genes are known in the art, for
15 example see J. Sutton and J. Hodgkin in 'The Nematode *Caenorhabditis elegans*' Ed. By William B. Wood and the Community of *C. elegans* Researchers CSHL, 1988 594-595; Zwaal et al; Target-Selected Gene Inactivation in *Caenorhabditis elegans* by using a Frozen Transposon
20 Insertion Mutant Bank' 1993, Proc. Natl. Acad. Sci. USA 90 pp 7431-7435; Fire et al, Potent and Specific Genetic Interference by Double-Stranded RNA in *Caenorhabditis elegans* 1998, Nature 391 860-811.

The possibility that *C. elegans* might be useful
25 for establishing links between compounds and specific *C. elegans* genes by virtue of comparison of phenotypes generated by exposure to particular compounds and by selected mutations is considered by Rand and Johnson in Methods of Cell Biology, Chapter 8, vol 84,
30 *Caenorhabditis elegans: Modern Biological Analysis of an Organism* Ed. Epstein and Shakes, Academic Press, 1995 and J. Ahringer in Curr. Op. in Gen. & Dev. 7; 1997; 410-415.

However, these authors observe and attribute
35 altered phenotypes on the basis of a single changed characteristic such as, for example, pharyngeal

- 5 -

pumping rate or defecation frequency. Since that single characteristic may be determined by expression of a number of genes and the operation of several biochemical pathways such a crude assessment of phenotype is not sufficient to establish a link between any one gene or pathway and a compound to which the worm has been exposed. As such the procedure would not be sensitive enough for resolution of the properties of thousands of compounds in a high throughput compound screen. An additional problem with the proposals of the prior art is that known phenotypic characteristics have all been described differently by different workers in the *C. elegans* field. Phenotype descriptions in the literature largely omit aspects not directly related to or not recognised to be related to the principle interest of the individual researcher. There is no standard nomenclature to identify a specific change. Without this it is impossible to equate newly observed phenotypes with particular known phenotypes for comparison purposes.

The present inventors have developed methods which solve these problems and thereby have converted *C. elegans* into a really useful tool in the drug discovery field. Specifically, in respect of each worm a 'phenotype profile' or 'fingerprint' is established based on looking for plurality of changed characteristics in a particular mutant or worm which has been exposed to an environmental change or a compound. Furthermore, each profile is scored by following a strict standard protocol of measurement and a standard description is applied to each characteristic. The determination of a phenotypic profile in this way for a plurality of mutants or worms exposed to compounds illuminates differences between different mutants or otherwise treated worms

- 6 -

which would not be apparent based on prior art methods. Furthermore, the standard scoring protocol and nomenclature allows the phenotypic profiles obtained to be collated into a library of reference profiles for direct comparison purposes. Thus, libraries of reference profiles can be established for mutant worms and for worms exposed to particular environmental changes or different sorts of compounds. Such libraries allow complex patterns of linkage to be established between particular compounds and particular genes or biochemical pathways and between individual compounds of known or unknown biochemical or pharmacological activity.

In accordance with a first aspect of the present invention there is provided a method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

- (a) providing a worm having a defect in at least one gene.
- (b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,
- (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotype profile associated with said defect,
- (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of worms each of which has a different defect, and
- (e) collating the phenotypic profiles so obtained into a library of said profiles.

- 7 -

Caenorhabditis elegans is the preferred nematode worm although the method could be carried out with other nematodes and in particular with other microscopic nematodes, preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size are extremely suited for use in mid- to high-throughput screening as they can easily be grown in the wells of a multi-well plate of the type generally used in the art to perform such screening.

It is preferred to establish the phenotypic profile on the basis of the measurement and scoring of at least three different characteristics, preferably at least six characteristics and more preferably at least ten characteristics. It will be appreciated that the more differences which can be scored between a worm with a genetic defect and a worm without the defect the better the resolution between different mutants. Although not limited to such, at least one of the plurality of changed characteristics which can be measured and scored may be selected from the list shown in Table 1, and possibly each of all the changed characteristics scored is one of those shown in Table 1.

In a preferred embodiment, the method used to establish the phenotypic profile comprises measurement and scoring of two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility. This list provides a core set of measurable characteristics which can be used to establish an informative phenotypic profile for any type of worm. Furthermore,

- 8 -

each of these characteristics is measurable using technical measuring apparatus, such as video image analysis, multiwell plate reader, and/or a technical assay procedure. In the most preferred embodiment, the method used to establish the phenotypic profile comprises measurement and scoring of all eight of the listed core characteristics. Measuring and scoring this set of core characteristics allows meaningful comparisons to be made between phenotypic profiles for worms subjected to diverse interventions. AS exemplified herein, comparisons can be drawn between profiles for two different mutant worms and between profiles for mutant worms and profiles for worms exposed to compound.

It is to be understood the terms "measuring" or "measurement" as used in connection with any of the methods described and claimed herein are to be interpreted as including not just absolute quantitative measurement wherein a numerical value is assigned to the characteristic but also comparative measurement, wherein characteristics of a worm which has been subject to an intervention (i.e. mutation, exposure to compound, exposure to environmental change) are measured relative to the same characteristics of a wild-type worm and scored as being 'larger', 'smaller', 'longer', 'shorter', 'fatter', 'thinner', 'darker', 'paler' etc.

For comparison purposes it is essential that the scored characteristics are represented in the same order for each profile. For standardization of procedure between different workers or to facilitate automation, measurement and scoring of the characteristics could be carried out in a pre-determined order according to a standard protocol. However, this is not essential to the operation of the method. In its simplest form and as shown in Example 5, the characteristics are recorded in a binary manner

- 9 -

as 'present' or 'not present' based on deviations from wild-type worms.

It is desirable to establish a library which comprises a phenotypic profile in respect of a defect in each gene in the worm genome and/or different defects in the same gene (allelic variations). As aforesaid there are a considerable number of available mutants (see Riddle, Blumenthal, Meyer and Priess and Ahringer above). In addition new ones can be generated by specific gene and site directed mutation and knock-out methods known to those skilled in the art such as ethyl methanesulphonate (EMS) mutagenesis, transposon insertion or genetic interference using double stranded RNA (see Sutton and Hodgkin, Zwaal et al and Fire et al above). The known or newly generated genetic defects may manifest themselves, for example, as the absence of expression of a gene, the reduction in expression of a gene, the over-expression of a gene, the expression of a functionally defective protein, the mis-expression of a protein, the ectopic mis-expression of a protein, the expression of a protein of altered stability or the alteration of gene expression as a function of time.

Generally, the manipulation of *C. elegans* to generate genetic defects can be carried out on wild-type worms or worms with existing single or multiple mutations. It may be desirable to genetically manipulate *C. elegans* carrying a reporter gene construct. The reporter molecule might be LacZ or green fluorescent protein but many other reporter molecules are known to those skilled in the art. Reporter gene constructs for *C. elegans* are described in Chalfie et al, 1994, Science 263 pp 802-805. It can also be desirable to genetically manipulate and then profile a transgenic worm, preferably a worm carrying a human gene, particularly where the gene is

- 10 -

associated with, or is a candidate for association with a human disease and therefore a putative drug target. A list of human diseases for which a particular gene has been implicated is given in the paper by J. Ahringer (see above) and also provided by OMIM. Center for Medical Genetics, John Hopkins University and National Biotechnology Information, National Library of Medicine, 1996. <http://www.ncbi.nlm.nih.gov/omim/>, although these lists are not necessarily exhaustive.

It is easy to establish transgenic lines in *C. elegans* and the methodology is described in Craig Mello and Andrew Fire, *Methods in Cell Biology*, Vol 48 Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

A form of the worm which may show a change in phenotype and may therefore be subject to profiling as described above is one in which the genetic defect and/or transgene and/or reporter gene is only present in a sub-set of the cells of the worm. It is possible for just the cells of a particular tissue to be the subject of a genetic manipulation.

The worm which is to be subject to determination of its phenotypic profile can be cultured by methods well-known in the art. *C. elegans* can grow on nutrient agar which has first been inoculated with bacteria on which the worms feed. Suitable culture methods are described in Rand and Johnson (see above) and in the examples given herein. Measurement of any changed characteristics which will determine the profile may be carried out using light microscopy, differential interference contrast optics or fluorescence microscopy. In addition immuno-chemical detection, colorimetric detection or detection of fluorescence, luminescence or radioactive labels may be used. In

- 11 -

some cases the changed characteristics may be biochemical only and might be detected, for example by a pH change in the growth media or a change in electrical potential. Different characteristics may need to be determined at different points in the growth cycle of the worm. For example, some phenotypic characteristics may be manifested only in the larvae while others are only detectable in the adult worm. In some cases it may be necessary to make several measurements of the same characteristic at pre-determined time intervals.

Phenotypic profiles generated by the methods described above can be collated into a library of profiles which are stored electronically on a database. However, it will be appreciated that the invention also provides a method of constructing a physical library or bank of worms each identifiable by their individual phenotypic profile. Such a worm library can be created using any or all of the methods described above and used for comparative purposes. The worms may be maintained by the culture methods described herein and/or frozen for long term storage by methods known to those skilled in the art. Libraries of phenotypic profiles or fingerprints of mutant worms or mutant worm libraries can be used to determine linkages between different genes and hence identify biochemical pathways. A particularly important use is the profiling of several mutations of the same gene and several genes of the same pathway. Different mutations in the same gene can have different phenotypes and often it is found that a careful analysis of the allelic series of a gene reveals important information that is hidden under a more severe phenotype of a null mutant (complete knock out, e.g. if it is lethal). Phenotypic profiles of different mutations of the same gene allow

- 12 -

characterisation of the gene by simply combining
(logical OR) the profiles of all the mutations,
whether they have been generated at the same time or
not. It is possible, however, to handle the mutations
separately and make more detailed connections, for
example, concerning protein domains in case the
similarity of phenotypes cluster with the sites of the
mutations.

Described above are methods for constructing a
library of phenotypic profiles for worms with a
plurality of genetic defects or a library of mutant
worms. However, in accordance with a second aspect the
present invention provides a method of constructing a
library of phenotypic profiles of nematode worms which
comprises the steps of:

- (a) exposing a worm to a compound,
- (b) measuring any changes in identifiable
characteristics of said worm as a result of
exposure to said compound,
- (c) systematically scoring a plurality of any
said changed characteristics to establish a
phenotypic profile associated with said
compound,
- (d) simultaneously or sequentially repeating
steps (a) to (c) in respect of each of a
plurality of different compounds, and
- (e) collating the phenotypic profiles so
obtained into a library of said profiles.

Methods for culturing *C. elegans* in the presence
of a test compound are described by Rand and Johnson

- 13 -

mentioned above and in the examples herein. In its simplest form a solution of the compound in a suitable solvent may be spread over a bacterial lawn on an agar plate before inoculation with the worm. Additional refinements include feeding the worm with bacteria, preferably *E. coli*, which have taken up the compound or attaching the compound to a carrier compound which is particularly attractive to the worm.

The worms which are exposed to the compound may be wild-type worms, mutant worms, transgenic worms and/or worms carrying reporter gene constructs as already described herein. Further the measurement and scoring of a plurality of changed characteristics is carried out by exactly the same procedures as already described herein for the phenotypic profiling of mutant worms. This must be a standard format in order that direct comparisons can be made between profiles obtained on exposure to compounds and profiles exhibited by mutants.

With compound screening it is possible to build up a series of different libraries depending on the compounds being tested. For example one library can comprise profiles generated in respect of each of the known compounds in a Pharmacopoeia, in other words compounds with known pharmacological activity.

Another library can comprise profiles generated by compounds known to interact with a particular biochemical pathway, which may or may overlap with those compounds from the Pharmacopoeia. Other libraries could include profiles for known compounds but with no known biological activity or compounds which are completely new molecules such as might be generated by combinatorial chemistry. As aforesaid the present invention is not limited to the production of phenotypic profile libraries but includes libraries or banks of worms whose phenotypic profile has been altered by exposure to compounds. In particular

- 14 -

embodiments assays may be carried out with several concentrations of the same compound, and/or with mixtures of compounds. For example compounds from compound libraries may each be tested individually or with one or more other influencing compounds. Furthermore, such compound testing protocols may be executed against identical worms or multiple mutant and/or transgenic backgrounds. In a particular example a panel of worm strains, covering a wide range of biochemical pathways and cellular activities by means of mutations in particular pathways, as well as reporter genes, is used for testing compounds. For each compound, potentially at several concentrations, a profile is recorded for the measurable phenotypes of each of the worm strains, either in parallel or sequentially.

In a third of its aspects the invention provides a method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

- (a) exposing a worm to an environmental change,
- (b) measuring any changes in identifiable characteristics as a result of said environmental change,
- (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said change,
- (d) simultaneously or sequentially repeating steps (a) to (c) for each of a plurality of different environmental changes, and

- 15 -

- (e) collating the phenotype profiles so obtained into a library of said profiles.

The environmental change may be, for example, a change in pH, osmolarity, temperature, exposure to radiation or exposure to bacteria or viruses. Each of these external influences may result in the manifestation of a different phenotypic profile of characteristics so that libraries of said profiles and affected worms can be constructed. Again, measurements and scoring of the profile should follow a standard protocol in order that valid comparisons can be made between these profiles and those in mutant and compound libraries.

The construction of worm and phenotypic profile libraries by the methods described above using the novel phenotypic profiling method described herein provides a very powerful tool for the discovery of new drugs. Profiles in each of the different libraries can be compared and links established between *C. elegans* genes and pathways, compounds and environmental effects. Preferably, the process of measuring and scoring the changed characteristics which go to make up the phenotypic profile is automated, making use of technical measuring apparatus. The profiles so generated may advantageously be stored electronically. Libraries of profiles can then be searched by computer which can identify identical or similar profiles, either within or between the different libraries. Quantitative data calculations, optionally in combination with boolean operations can be used.

A comparison of the profile generated by a particular compound with the profiles of particular mutants may indicate the likely gene or biochemical pathway with which the compound interacts in the worm. Other databases can then be searched for a match of

- 16 -

the worm gene with an equivalent human gene. The human gene might already be associated with a human disease as could be determined for example, from the OMIM database mentioned above. Thus, by use of the worm screen a potential candidate drug can be identified. The discovery of the mode of action of a compound with known pharmacological or biochemical activity is facilitated by comparing its phenotypic profile in the worm with the mutant library or environmental change library of profiles to identify possible targets for the compound. other possibilities include finding a new potential medical indication of a known compound, a medical indication for a novel compound, an alternative method of treatment of a known disease or an indication of the reason for the side effect exhibited by some known pharmaceuticals. Testing worms with compounds, scoring the phenotypic profile in the novel manner described herein and then searching previously established libraries of profiles can potentially achieve all those goals. Once a compound has been identified as having the potential to be a therapeutic agent it can be processed through the more traditional drug discovery routes. The compound can be tested in more specific in vitro tests based on the new knowledge of the target for the compound and in animal models of the target disease. Structural variants then can be generated by medicinal chemistry with a view to improving activity.

The invention will now be described with reference to the following Examples, together with accompanying Figures, in which:

Figure 1 is a schematic diagram of the left lateral view of the body of *C. elegans*. The body of *C. elegans* is divided into a head, a body and a tail region. The head region stops at the end of the

- 17 -

pharynx, the body stops at the rectum and the tail includes the tail whipe. *C. elegans* usually crawl on the right side. The ventral located vulva defines the ventral side of *C. elegans*.

5 Figure 2 is a schematic diagram of *C. elegans* showing the characteristics "hypertrophy of the head and "extensions on head".

Example 1

10 General Profiling by Plate Drop Assay

4ml NGM agar (see 'The Nematode *Caenorhabditis*
Elegans' Ed. by William B. Wood and the Community of
C. elegans Researchers CSHL, 1988, pg 589) is poured
15 into 3cm plate, and seeded with approximately 5µl of
an *E. coli* overnight culture and grown preferably for
one week at room temperature. If a compound is to be
profiled 10µl of compound dissolved in DMSO or other
appropriate solution is pipetted onto the bacterial
20 lawn. The lawn should be covered completely. (This
step can be omitted if a mutant, transgenic or other
worm is being profiled without compound). After
overnight soaking in of compound one *C. elegans* (L4
stage) per plate is put in the bacterial lawn. Worms
25 are checked after some hours, plates are incubated at
21°C and worms screened for phenotypes (control have
L1 progeny growing). Plates are checked again after 4
days for phenotypes of F1 progeny (control shows all
stages up to gravid hermaphrodites). Plates which have
30 to be looked at again on subsequent days because of
slow growth or for further checks are put aside. A
plate protocol sheet such as that shown in Table 2 is
completed deciding on one of the following routes: no
effect/unspecific effect/needs to be applied at lower
35 concentrations/needs to be profiled. If concentrations
are appropriate and a decision can be made scoring of

characteristics to produce a profile can be started using the profiling list in Table 1. Because the compound is pipetted onto a bacterial lawn rather than it being incorporated into the agar, as has been done in the prior art, this method is designated a 'plate drop assay'.

Table 1

1. Compound specific phenotypes

[illegible]

2. Viability

[illegible]

- 19 -

2.3 Embryonic arrest f F1									
2.3.1 Leakyn ss									
2.3.2 Appearance of eggs									
2.3.2.1 dark eggs									
2.3.2.2 bright eggs									
2.3.2.3 two-fold or older									
2.3.2.4 irregular egg size									
2.4 Larval arrest of F1									
2.4.1 Leakyness									
2.4.2 at L1									
2.4.3 at L2									
2.4.4 at L3									
2.4.5 at L4									
2.5 Embryonic arrest of F2									
2.5.1 Leakyness									
2.5.2 Appearance of eggs									
2.5.2.1 irregular egg size									
2.6 Larval arrest of F2									
2.6.1 Leakyness									
2.6.2 at L1									
2.7 Died during adulthood (F1)									
2.8 Died during adulthood (F2)									

3. Life cycle

Phenotype									Comment
Abnormal									
3.1 Growth abnormal									
3.1.1 only generation cycle slowed down									
3.1.1.1 oldest stage L1									
3.1.1.2 oldest stage L2									
3.1.1.3 oldest stage L3									
3.1.1.4 oldest stage L4									
3.1.2 generation cycle slowed down while displaying defects									
3.1.2.1 oldest stage L1									
3.1.2.2 oldest stage L2									
3.1.2.3 oldest stage L3									
3.1.2.4 oldest stage L4									
3.1.3 stage changed									
3.1.3.1 delayed hatching									
3.1.3.2 arrested growth in L1									
3.2 Dauer formation defective									
3.2.1 constitutive dauer									
3.2.2 non-conditional constitutive									
3.2.3 defective									
3.2.4 dies on recovery									
3.3 Life span changed									
3.3.1 Life span is shorter									
3.3.2 Life span is prolonged									

4.5 Cuticl defects									
4.5.1 blistered									
4.5.1.1 symmetrically									
4.5.1.2 around the head									
4.5.1.3 around the pharynx									
4.5.1.4 around the body									
4.5.1.5 around the tail									
4.5.2 moulting defective									
4.5.2.1 incomplete molts									
4.5.2.2 supernumerary molts									
4.5.3 burst -									
4.6 Poured out									

5. Movement

[illegible]

[illegible]

6. Mechanotransduction (Touch with a wire and with eyelash)

[illegible]

- 23 -

7. Sensory system

Phenotype										Comment
Abnormal										
7.1 Avoidance of bacteria										
7.2 Bordering behaviour										
7.3 Chemotaxis defective										
7.3.1 attraction										
7.3.2 avoidance										
7.4 Thermotaxis defective										
7.4.1 attraction										
7.4.2 avoidance										

8. Environmental response

Phenotype										Comment
Abnormal										
8.1 Osmolarity sensitive										
8.2 Thermotolerance changed										
8.3 UV Resistance changed										
8.4 Oxygen sensitive										

9. Pharynx

Phenotype										Comment
Abnormal										
9.1 Pharynx stuffed										
9.2 Morphology defects										
9.3 Pumping defects										
9.3.1 pumping reduced										
9.3.2 pumping enhanced										
9.3.3 pumping irregular										
9.3.4 no pumping										
9.4 Eating defective										

10. Intestine

Phenotype										Comment
Abnormal										
10.1 Morphology defects										
10.1.1 enlarged										
10.1.2 detached										
10.2 Color of contents										
10.2.1 darker										
10.2.2 lighter										

- 24 -

11. Rectum

Phenotype										Comment
Abnormal										
11.1 Morphology defects										
11.1.1 protruding										
11.1.2 scarring										
11.1.3 absent										
11.2 Constipation										
11.2.1 foregut filled/enlarged										
11.2.2 hindgut weak										
11.2.3 hindgut strong										
11.3 Defecation cycle defective										
11.3.1 expulsion defective										
11.3.1.1 weak expulsion										
11.3.1.2 no expulsion										
11.3.2 aBoc defective										
11.3.3 pBoc defective										
11.3.4 wrong timing of cycle										

12. Gonad

Phenotype										Comment
Abnormal										
12.1 Morphology defects										
12.1.1 defective gonad										
12.1.2 one arm missing										
12.1.3 multiple gonad										
12.1.4 monopolar gonad forward										
12.1.5 monopolar gonad backward										
12.1.6 no gonad										
12.2 Light brown										

13. Vulva

Phenotype										Comment
Abnormal										
13.1 Morphology defects										
13.1.1 defective vulva										
13.1.2 protruding vulva										
13.1.3 multi vulva (number)										
13.1.4 no vulva										
13.1.5 leaky vulva										
13.1.6										
13.1.7										

14. Fertility

Phenotype										Comment
Abnormal										
14.1 Brood size abnormal										
14.1.1 small r										
14.1.2 larg r										
14.2 Egg laying defect										
14.2.1 no gg retention										
14.2.2 immediate Egl										
14.2.3 progressive Egl										

- 25 -

14.2.4	egg laying defective								
14.2.4.1	weak Egl								
14.2.4.2	strong Egl								
14.2.5	bloated worms								
14.2.5.1	weak bloating								
14.2.5.2	strong bloating								
14.2.5.3	bags of worms								
14.2.6	no egg laying								
14.3	Only oocytes								
14.4	Sterile								
14.5	Maternal effect sterile								

15. Male

Phenotype									Comment
Abnormal									
15.1 Frequency									
15.1.1	high incidence of males								
15.2	Mating defective								
15.3	Morphology								
15.3.1	leptoderan tail								
15.3.2	scrawny								
15.3.3	copulatory plug								
15.4	Mating behaviour								
15.4.1	defective sensory contact								
15.4.1.1	no response to dorsal contact								
15.4.1.2	no response to ventral contact								
15.4.2	defective backing								
15.4.2.1	no backing								
15.4.2.2	no continued backing								
15.4.3	defective turning								
15.4.3.1	loose turns								
15.4.3.2	stop at the tail								
15.4.3.3	slide off the tail								
15.4.4	defective vulval location								
15.4.5	defective spicule insertion								

16. Progression of phenotype

Phenotype									Comment
Abnormal									
16.1	Dependent on generation								
16.1.1	F1 different from P0								
16.1.1.1	weaker								
16.1.1.2	worse								
16.1.1.3	lower penetrance								
16.1.1.4	higher penetrance								
16.1.1.5	not affected								
16.1.2	F1 different from F2								
16.2	Dependent on stage								
16.2.1	appearance of phenotype								
16.2.1.1	after L2								
16.2.1.2	during adulthood								
16.2.2	shift of phenotype								
16.3	Dependent on age								
16.3.1	phenotype gets worse								
16.3.2	phenotype gets better								

Tabl 2

plat	well	by	date
negative control	positive control	finished	confirmed (≥ 3 worms)
no effect	unspecific effect	needs to be applied at lower concentrations	needs to be profiled

Day 0

compound
invisible
coloured
droplets
crystals
complete crust

bacteria
normal lawn
grown as ring
thin
crust
died

worm
happy
run away
irregular movement
slow movement
no movement

Day 1

appearance
healthy
slightly unhealthy
slightly starved
strong starved
very sick

worm gone
lost
suicide
in agar
starved outside
died in compound

replaced by
number and stage
left progeny

movement
normal
tracks more outside
tracks not in center
amplitude increased loopy
amplitude variable
amplitude decreased
enhanced movement
slow movement
no movement
specific

body
normal gravid adult
pumping defects
light brown messy gonad
pale with dark spots
few eggs in gonad
pharynx stuffed
foregut filled large
hindgut constipated
protruding vulva
other:

progeny
normal
reduced broodsize
younger staged
oocytes
coagulated eggs
dead eggs
dying hatchlings
crippled larvae

Day 4

food
still plenty of
already finished
finished soon
outside comp.
not eatable, died

adult viability
still fertile
laying oocytes
died
died as bag of worms
missing

growth rate
normal
reduced broodsize
younger staged

movement
normal
population more outside
population not in center
amplitude increase, loopy
amplitude variable
amplitude decreased
enhanced movement
slow movement
no movement
specific:

body
normal gravid adult
pumping defects
light brown messy gonad
pale with dark spots
few eggs in gonad
pharynx stuffed
foregut filled large
hindgut constipated
protruding vulva
other:

brood viability
dead eggs
dead larvae
larval arrest
later scoring
day of screen
day of worm

comparison of phenotypes

progeny shows PC phenotype
similar
worse
a few only
weaker
no effect

new worms show phenotype
similar
worse
not all
weaker
not effect

stage & age
all stages
young only
late larvae and adults
adults only
old adults

comparison to other plates

comparison to known drugs

comparison to known mutants

- 27 -

Example 2**Profiling of a compound library (new compounds)**

5 To profile new compounds from a library, the general
profiling protocol is followed with the variations.
Compounds are profiled once in undiluted
concentration, the actual concentration being
dependent on the compound library in question but will
be between 0.01 mg and 1 mg of compound/10 μ l DMSO.

10

For compounds with a MW of 500 this calculates to 2-
200 mM stock. Dilution in 4ml agar would be at 5-500
 μ M. The high dose may create lots of unspecific effect
problems e.g. bacterial death and worm starvation.

15

Thus, if necessary the compounds are applied in a
second round at lower concentrations which are
dilutions in DMSO of 1/3, 1/10 and 1/30 of the
undiluted concentration. A concentration is finally
chosen for each compound which will allow a phenotype
20 profile to be established according to the standard
procedure.

20

Example 3**Profiling of known compounds (biotools, pharmacopoeia)**

25

To profile known compounds from a library the general
profiling protocol is followed with the following
variations. The stock solution is preferred as 100mM
in DMSO and the experiment is started *ab initio* with a
concentration series. The concentration series is used
30 as described below. In one series of concentrations 15
or so worms (for a reasonable number of short term
effects) are placed in the agar. In three series 1
worm each is placed on the agar to score a reasonable
35 number of progeny. Lost worms of the latter three
series of concentrations can be replaced from the

30

35

- 28 -

large pool where worms have been exposed to the compound in the same way. The following concentrations can be used:

5	conc.in 10 μ l drop	100mM	30mM	10mM	3 mM	1mM	0.3mM
	conc.in 4ml drop	100 μ M	300 μ M	100 μ M	30 μ M	10 μ M	3 μ M

Example 4

10 Comparison of agar assay to drop assay

A set of compounds from the pharmacopoeia have been profiled using the general protocol (all compounds were of known activity and are described in
15 Martindale: The Complete Drug Reference, 32nd edition, Pharmaceutical Press 1999). The plate drop assay was compared against standard of pouring compounds into the agar as described in literature which method is designated agar assay. In the drop assay as well as in
20 the agar assay, the compounds were added to the worm in a variety of concentrations, and the survival of the worm was scored as well as the phenotypic profile induced by the compound. The lowest concentration of a compound, still resulting in the death of the nematode
25 was designated minimal lethal dose. The maximal concentration of a compound that did not result in the death of the nematode was designated maximal nonlethal dose. The minimal concentration of a compound that still resulted in a measurable phenotype was
30 designated minimal effective dose. The concentrations of the compounds in the agar assay were compared to the concentrations in the drop assay. From this observation one may conclude that the newly described drop assay protocol turns out to be far more efficient
35 for most compounds. The following table lists the calculated concentration ratio needed to get the same

- 29 -

effect with the compound in the agar assay (in 2 ml agar) rather than the drop assay (in 4 ml agar).

Table 3:

Compound	Site	min. lethal dose	max. nonlethal dose	min. effective dose	average potency ratio
ketanserine	serotonin rec. agonist	>610			610
tamoxifen	estrogen rec. antagonist	204	304		254
fluoxetine	serotonin reuptake inh.	124	186		154
pancuronium	nicotinic antagonist			>100	100
methoxyphenylpiperazin	α -adrenorec. ligand	>48	>146	72	88
naloxone	opioid antagonist		>44	78	60
diheptylbipyridinium	ryanodine rec. antag.	20	30	36	28
W7	calmoduline antag.	20		10	14
thapsigargin	serca antagonist				14
physostigmine	cholinesterase inh.			8	8
lobeline	nicotinic rec. ligand			4	4
riluzole	glutamase release inh.	2	2	4	2
levamisole	acetylch. rec. antag			$\frac{1}{2}$	$\frac{1}{2}$
nicotine	acetylch. rec. antag			$\frac{1}{2}$	$\frac{1}{2}$

Minimal lethal dose: rate between the lowest concentration in which the compound is lethal to the worm in both assays
 Maximal non-lethal dose: rate between the highest concentration in which the compound is not lethal in both assays
 Minimal effective dose: rate between the lowest concentration in which the compounds results in a phenotype in both assays
 Average: average of the rates

- 30 -

Example 5**Preferred set of informative characteristics**

5 Worms exposed to a compound, carrying a mutation or
are transgenic are examined for the following 8
informative features/phenotypes:

1. Viability

10 Worms are examined for viability at all stages of the
life cycle, being embryogenesis, larval stages 1 to 4
and adulthood. Dead embryos are defined by not
hatching within 24h and dead worms are defined by not
moving, by lack of pharynx pumping, by sick or pale
appearance and by lack of response to mechanical
15 stimulation.

Method:

Embryonic lethality is measured by counting the amount
of unhatched worms after 24 hours (Elispot, Zeiss).
20 Counting of unhatched worms could also be automated
using the FANS device, described below. Viability of
larvae and adults is measured by dye uptake.

2. Life cycle

25 Progeny are examined for the length of the generation
cycle in comparison to control progeny (of a wild-type
worm). The stage of a synchronized progeny will be
compared to the stage of a synchronized control
progeny (N2, Bristol strain) after three days at 20°C.
30 The developmental stages can be distinguished by vulva
development, expression of stage-specific markers,
such as collagen IV, body length and transparency.

Method:

35 Measuring the body length of a population allows
determination of the actual stage in the life cycle

- 31 -

(For body shape measurement, see 3. Body shape).

Expression of stage-specific markers can be examined using antibodies of the appropriate specificity, by way of example an antibody that recognizes an antigen on the surface of *C. elegans* L1 larvae has been described by Hemmer *et al.*, (1991) *J Cell Biol*, 115(5): 1237-47.

3. Body shape

Worm size is determined by measuring worm length and worm diameter.

Method:

The body length of a synchronized progeny of adult worms is compared to the body length of a synchronized control progeny (N2, Bristol strain). Measurement of body length can be achieved using a 'worm dispenser apparatus' which is commercially available from Union Biometrica, Inc, Somerville, MA, USA. This apparatus has properties analogous to flow cytometers, such as fluorescence activated cell scanning and sorting devices (FACS). Accordingly, it may be commonly referred to as a "FANS" apparatus, for fluorescence activated nematode scanning and sorting device (FANS). The FANS device enables the measurement of properties of microscopic nematodes, such as size, optical density, fluorescence, and luminescence.

Body size may also be measured via image analysis, in which case the measurements recorded may include worm diameter and deviation from the typical tube shape of a wild-type worm.

4. Movement behaviour

The measurement of movement behaviour can include measurement of the speed of movement, or of the

- 32 -

pattern of movement (e.g. direction) or both. A wild-type worm moves in a sinusoidal way forward and pauses or moves backward occasionally. Any deviation from this wild-type pattern of movement can be scored as a 'changed' characteristic.

Method:

An assay based on the following principles may be used to determine the speed of movement of a worm culture:

Nematode worms that are placed in liquid culture will move in such a way that they maintain a more or less even (or homogeneous) distribution throughout the culture. Nematode worms that are defective in movement will precipitate to the bottom in liquid culture. Due to this characteristic of nematode worms as result of their movement phenotype, it is possible to monitor and detect the difference between nematode worms that move and nematodes that do not move.

Advanced multi-well plate readers are able to detect sub-regions of the wells of multi-well plates. By using these plate readers it is possible to take measurements in selected areas of the surface of the wells of the multi-well plates. If the area of measurement is centralized, so that only the middle of the well is measured, a difference in nematode autofluorescence (fluorescence which occurs in the absence of any external marker molecule) can be observed in the wells containing nematodes that move normally as compared to wells containing nematodes that are defective for movement. For the wells containing the nematodes that move normally, a low level of autofluorescence will be observed, whilst a high level of autofluorescence can be observed in the wells that contain the nematodes that are defective in movement.

- 33 -

In an adaptation of the movement assay, autofluorescence measurements can be taken in two areas of the surface of the well, one measurement in the centre of the well, and one measurement on the edge of the well. Comparing the two measurements gives analogous results as in the case if only the centre of the well is measured but the additional measurement of the edge of the well results in an extra control and somewhat more distinct results.

As an alternative to the above-described movement assay, specialist software such as SIMI Scout (designed for movement study of an athlete) may be used to determine speed of movement, deviation from sinusoidal movement and even the overall pattern of movement of the worm.

5. Mechanotransduction

Worms are examined for response to mechanical stimulation.

Method:

When the plate on which *C. elegans* are cultured is dropped wild-type worms react by enhanced movement and enhanced overall activity. The capability of a worm to respond to a mechanical stimulus is measured by the difference in speed of movement before and after stimulation.

6. Pharynx pumping

The phenotypes "Pumping frequency reduced, Pharynx pumping irregular" etc. describe the activity of the cyclic contraction of the pharynx muscles that occurs in a feeding adult about 3 times in a second. The contraction cycle can be described as the nearly simultaneously contraction of the corpus, anterior

- 34 -

isthmus, and terminal bulb, followed by relaxation.

Method:

5 The following pharynx pumping characteristics may be
analyzed by image analysis: The frequency of pumping
by counting the pharynx contraction. Pharynx
contraction can be measured visibly by the opening and
closing of the anterior corpus. The time of opened
10 anterior corpus and the diameter of the opened corpus
is used to measure hypercontraction, relaxation and
strength of a contraction.

15 The following is an example of a pumping assay which
allows measurement of the total efficiency of feeding
of a worm, which is related to pumping:

20 The pumping rate of the pharynx is measured indirectly
by adding a marker molecule precursor such as calcein-
AM to the medium and measuring the formation of marker
dye in the *C. elegans* gut. Calcein-AM is cleaved by
esterases present in the *C. elegans* gut to release
calcein, which is a fluorescent molecule. The pumping
rate of the pharynx will determine how much medium
will enter the gut of the worm, and hence how much
25 calcein-AM will enter the gut of the worm. Therefore
by measuring the accumulation of calcein in the
nematode gut, detectable by fluorescence, it is
possible to determine the pumping rate of the pharynx.

30 To perform the pharynx pumping screen with calcein-AM,
a concentration of between 1 and 100 μ M calcein-AM is
added into the medium. Preferably 5 to 10 μ M calcein-
AM is used. Fluorescence is measured using a multi-
well plate reader (Victor2, Wallac Oy, Finland) with
35 following settings: Ex/Em = 485/530.

- 35 -

7. Defecation

The defecation of *C. elegans* is a recurrent event comprising of the following steps: pBoc, aBoc and expulsion. Defecation in nematodes such as *C. elegans* is achieved by periodically activating a defined sequence of muscle contractions. These contractions are started in the anterior body wall muscles. At the zenith of the anterior body contractions the four anal muscles also contract. The four anal or enteric muscles are the two intestinal muscles, the anal depressor and the anal sphincter. In addition to this series of muscle contractions, specific neurons are also involved in the regulation of defecation, including the motor neurons, AVL and DVB.

Method:

In order to construct a phenotypic profile, well-fed adults are typically examined after one day for constipation. The time between two pBocs is also scored.

The rate of defecation of *C. elegans* can also be quantitatively measured using an assay based on the following principles:

The rate of defecation of nematodes such as *C. elegans* can be easily measured using a marker molecule which is sensitive to pH, for example the fluorescent marker BCECF. This marker molecule can be loaded into the *C. elegans* gut in the form of the precursor BCECF-AM which itself is not fluorescent. If BCECF-AM is added to nematode culture medium in the wells of a multi-well plate the worms will take up the compound which is then cleaved by the esterases present in the *C. elegans* gut to release BCECF. BCECF fluorescence is sensitive to pH and under the relatively low pH

- 36 -

conditions in the gut of *C. elegans* (pH<6) the compound exhibits no or very low fluorescence. As a result of the defecation process the BCECF is expelled into the medium which has a higher pH than the *C. elegans* gut and the BCECF is therefore fluorescent. The level of BCECF fluorescence in the medium (measured using a multi-well plate reader on settings Ex/Em=485/550) is therefore an indicator of the rate of defecation of the nematodes.

8. Fertility

A wild-type adult hermaphrodite *C. elegans* lays about 8 eggs per hour.

Method:

The amount of eggs laid by 20 hermaphrodite *C. elegans* during at least 60 min is counted. The amount of eggs may be counted by simple visual inspection or using a FANS device, described above.

Example 6

Comparison of profiles within a library

(daf-4 belongs to two pathways)


Mutant worms have been profiled according to the general profile protocol. Table 4 shows a summary of the profile, also called fingerprints, of one mutation of the indicated genes. Entries are binary with empty fields indicating a phenotype (deviation from negative control, here wild-type) not found assuming that it could have been measured. Any other entry including comments or quantitative data is read as measured phenotype in this binary scheme and indicated by *. The table lists only phenotypes that do have a

- 37 -

positive entry, not necessarily complete, leaving pages of empty fields alongside and arranged according to a particular enquiry. The upper half consists of the hierarchical categories "dauer formation phenotypes" and "body shape phenotypes" as well as their relevant sub-phenotypes. The lower part consists of a set of hierarchically unrelated phenotypes subsumed under the enquiry categories, "increased activity" and "decreased activity". The complete list of characteristics is to be found in Table 1.

The point of including the lower part is to show the principle of recording all observed phenotypes, that they can be used to distinguish similar phenotypic profiles in detail and that they can be arranged in order to make comparisons. In this case it is seen that the dichotomy of long versus short body length does not correlate to the dichotomy of increased versus decreased activity.

The upper part shows 5 genes (i.e. a mutation in that gene) affecting dauer formation as well as 5 genes affecting body shape in a particular combination. A mutation in one gene, daf-4, is unique in sharing the characteristics of both phenotypic groups. The following picture illustrates the phenotypic overlap as found by comparing entries in the phenotypic profiles.



From this overlap a hypothesis of a mechanistic link can be put forward for daf-4. In this particular case the mechanistic link is confirmed by the molecular

- 38 -

nature of the genes, which as far as known are all members of the TGF β pathway by sequence similarity:

		dbl-1 TGF β like ligand
5	daf-7 TGF β like ligand	sma-6 type I receptor
	daf-1 type I receptor	daf-4 type II receptor
	daf-4 type II receptor	sma-2 SMAD
	daf-3 SMAD	sma-3 SMAD
	daf-14 SMAD	sma-4 SMAD

The DAF-4 protein probably acts as a type II receptor in both pathways. The similarity of phenotypic profiles allows one to hypothesize mechanistic relationships in a manner analogous to sequence similarity of genes. For example a compound which induces the phenotypes: longer or shorter body length in combination with 2 or 3 of pale, thin and variable egg size, in worms exposed to it, is very likely to act on a protein of the TGF β pathway.

Table 4:

Phenotype	<i>daf-1</i>	<i>daf-7</i>	<i>daf-3</i>	<i>daf-14</i>	<i>daf-4</i> <i>e1364</i>	<i>sma-2</i> <i>e502</i>	<i>sma-3</i> <i>e491</i>	<i>sma-4</i> <i>e729</i>	<i>lon-1</i> <i>e185</i>	<i>lon-3</i> <i>e2175</i>
25 dauer formation	●	●	●	●	●					
constitutive dauer					
recovery defective					
body shape					●	●	●	●	●	●
30 short						
long									.	.
thin				
pale					

- 39 -

Phenotype	<i>daf-1</i>	<i>daf-7</i>	<i>daf-3</i>	<i>daf-14</i>	<i>daf-4</i> <i>e1364</i>	<i>sma-2</i> <i>e502</i>	<i>sma-3</i> <i>e491</i>	<i>sma-4</i> <i>e729</i>	<i>lon-1</i> <i>e185</i>	<i>lon-3</i> <i>e2175</i>
irregular egg size				
increased activity					●		●	●	●	●
enhanced movement					.		.		.	
5 amplitude increased									.	
head movement enhanced						
foraging behaviour increased					.			.		.
pharynx pumping enhanced							.		.	
constitutive pumping							.	.	.	
10 no egg retention									.	.
decreased activity						●				
lay still						.				
slow movement						.				
15 pharyngeal pumping reduced						.				

Example 7

20 **Comparison of phenotypes induced by acetylcholine
esterase inhibitors**

Wild type *C. elegans* adults have been exposed to
acetylcholine esterase inhibitors at various
25 concentrations. The worms have been profiled over two
generations, meaning four profiles have been
generated. All phenotypes from the phenotype list are
displayed that have been measured in this experiment.
Two phenotypes "loopy head movement" and "body dragged
30 by head" are shared by most of the esterase
inhibitors. This is called phenotype activity

- 40 -

relationship (PAR, by analogy to structure activity relationship SAR). The shared phenotypes are used to identify the action of a new compound. The unshared phenotypes are used to distinguish drugs or unravel side effects when these phenotypes are part of another PAR.

Table 5:

Phenotypes	Physostigmine	Neostigmine	Ambenonium	Tacrine	Galantamine	Trichlorfon
Thin	X					
Lay still	X					
Erratic	X					
Weak kinker		X				
Jerky				X		X
Enhanced head movement						X
Loopy head movement	X	X		X(L1)		X
Body dragged by head	X	X				X
Irregular touch response	X	X				
Reduced brood size	(X)					X
Delayed growth						X

Example 8

Comparison of phenotypes of mutations in the acetylcholine neurotransmission pathway

C. elegans adults and larval stages that are homozygous for the mutations *cha-1*, *unc-17*, *snt-1* and *cat-1* have been profiled, meaning fingerprints have been generated. All phenotypes from the phenotype list are displayed that have been scored in this

- 41 -

experiment. The phenotypes "small", "resistance to CHA inhibitors (Ric)", "slow pumping" and "slow growth" are shared. This is called phenotype activity relationship (PAR, in analogy to structure activity relationship SAR). The shared phenotypes are used to identify genes in a pathway. The unshared phenotypes are used to distinguish these genes or unravel further functions in parallel or new pathways when these phenotypes are part of another PAR. The fingerprint of *cat-1* is different because this gene is involved in the dopamine pathway.

Table 6:

Phenotype	<u><i>cha-1</i></u> ChAT (synthesis)	<u><i>unc-17</i></u> VchAT (ACh- transporter)	<u><i>snt-1=ric-2</i></u> Synaptotag min homolog	<u><i>cat-1</i></u> VMAT (monamine- transporter)
Coiler	X	X		
Small	X	X	X	
Slow growth	X	X	X	
Ric	X	X	X	
Slow pumping	X	X	X	
Jerky when backing	X			X
Low ChAT level	X			
Poor male turning				
Enhanced foraging behaviour				
Enhanced foraging behaviour				
Defecation defects				
Shrinker-uncs				

- 42 -

Example 9**Method to profile an intervention (mutation, compound etc)**

- 5 Profiling a mutation in the gene *unc-17* that affects transportation of acetylcholine.

10 In the literature this phenotype is described, concerning movement, body size and feeding, as severe coiler, being rather small and thin and has only slow, irregular pumping of the pharynx (Riddle et al., "C. elegans II" Cold Spring Harbor Laboratory Press, 1997). By systematically describing *unc-17* the

15 resulting fingerprint unravels more details and new properties: Concerning movement, body size and feeding the phenotypes strong coiler, spiralling inwards posteriorly, curly jerky and moves better forward, being small have been profiled. In addition defects in the sensory system, defecation and reproductive system

20 have been found, in detail: the touch response is gone, constipation, aberrant defecation cycle (aBoc) and egg laying defective (no egg retention).

25 **Example 10**

Method to add biological information to a particular phenotype

30 One phenotype of the mutation *unc-4* is "coiler" (looks like a snail). The fingerprint of *unc-4* adds for "coiler" the details "ventral side out" and "spiralling inwards posteriorly". This occurs when a set of neurons that control the forward movement of the ventral part of the worm (VA2 - VA10) gets the

35 same input than another set of neurons that controls the backward movement of the ventral part (VB2 -

- 43 -

VB10).

In this case the ventral muscles get contradicting signals and only the dorsal muscles contract properly. The result is a coiler that has only the ventral side outwards. We explain most of the phenotypes as consequence of a mislead process, here synaptic input.

10 **Example 11**

Comparison of phenotypes induced by compounds acting on GABAnergic neurotransmission

15 Wild-type *C. elegans* adults have been exposed to GABA agonists (Muscimol) and GABA antagonists (Ivermectin and Fipronil) at various concentrations. Worms have been profiled and the scored phenotypes are displayed as fingerprints.

20 In addition, two mutations in the GABAnergic pathway have been profiled and compared with the compound induced phenotypes: *unc-25* encodes for the decarboxylase and *unc-49* encodes for a GABA receptor.

25 The phenotype "shrinker" is present in all fingerprints (see Table dark grey). This phenotype is used as marker or diagnostic phenotype to identify activity of a compound or gene in the GABAnergic pathway. There are further phenotypes only shared by
30 some compounds and mutants (see Table light grey). These phenotypes are used to build a phenotype activity relationship (PAR).

35 The shared phenotypes are used to identify the action of a new compound when "shrinker" cannot be used or to reveal more details on a compound action. For example,

- 44 -

all compounds and *unc-25* fingerprints contain constipation phenotypes but not the fingerprint of *unc-49*, although GABA is used for the defecation process. This is coincident with earlier findings that the UNC-49 gene product is not required for defecation.

These results may indicate the existence of another yet unknown GABA receptor in *C. elegans*. The unshared phenotypes are used to unravel toxic side effects or other mode of actions.

Table 7:

Phenotypes	Muscimol	Ivermectin	Fipronil	<i>unc-25</i>	<i>unc-49</i>
Pale	x	x		X	
Motionless (paralyzed) I	x	x			
Nearly motionless	x	x			
No movement but motion II	x		x	X	x
Little movement	x		x	X	x
Slow movement III	x		x	X	
Enhanced movement V					
Stiff rods					
Loose rods	x	x			
Rigid paralysis (hypercontracted)					
Flaccid paralysis (relaxed)	x	x			
Bent body, jerky body, abnormal Omega appearance			x	(x)	
Enhanced foraging			x		x
Shrinker before movement	x		x	X	
Shrinker	x	x	x	X	x
No pumping	x	x			
Weak pumping					
Pumping frequency reduced		x	x		
Pumping frequency enhanced					
Pumping irregular	x				
Constipation		x	x	X	
Foregut filled/enlarged			x		
Hindgut weak constipated		x	x	X	
Hindgut strong constipated				X	
Defecation cycle defective	x	x	x	x	
(time: pBoc)					
Weak expulsion				x	
No expulsion				x	
No egg retention (12-cell stage)					
Weak egg laying defect (comma)					
Strong egg laying defect (pretzel)		x	x		
Blocked worms			x		
Bags of worms			x		

Example 12**Definition of body shape phenotypes**

Aberrations of the body shape of *C. elegans* can be the result of mutations in a vast amount of genes. These genes may be required directly for the formation of the hypodermis, the hydroskeleton and the correct patterning of the worm body plan, e.g., collagen or even-skipped. They could be involved in the control of growth or metabolism like genes of the TGF β pathway or genes required for feeding. Eventually, mutations in certain genes that cause primary defects, e.g., absence of head muscle, cause secondary defects in the body shape like dystrophy in the head region.

Body shape phenotypes are all visible or measurable deviations of the body shape, colour and content. Phenotypes are comparatively measured against wild-type (N2, Bristol strain) and scored as deviation of wild type in the corresponding developmental stage, sex and preparation. The scored phenotype comes with the percentage of worms positive for that phenotype within a population.

Table 8: Scientific definition of body shape phenotypes. The phenotypes listed in the left column are described and defined in the right column. Some phenotypes are derived from the classical worm jargon like "dumpy", which is still shorter than "short and thick worm".

PHENOTYPE	DEFINITION
Proportion abnormal	
Short	Body length less than wild type.
Long	Body length more than wild type.
Thin	Body diameter less than wild type.
Thick	Body diameter more than wild type.

- 46 -

Dumpy	Body length less but body diameter more than wild type.
Spindle-shaped	body diameter is more for only a restricted region of the body.

Head defects

Hypertrophy of the head	Regions of the head are thickened. This additional tissue is part of the head and enclosed by the hypodermis.
Extensions of head	Small hypertrophied regions of the head.
Notched head	Extensions, protrusions on the dorsal side of the head.
Hammer head	Extensions at the head tip resemble a hammer like appearance.
Dystrophy of the head	Regions of the head are thinned due to missing tissue.
Swollen	The head looks like a balloon.
Rounded	The tip of the head is rounded.
Tapering	The tip of the head is tapering.
Vacuoles only in head	Vacuoles visible in the head but not in the rest of the body.
Only head bent	The head is held most of the time in a bent position. In extreme cases the worm looks like a walking stick.
Autodecapitation	The head/body connection is thinner, which results occasionally in an autodecapitation due to a body wall muscle contraction.

Body defects

Scrawny	Worm is shorter, thinner, pale and sick.
Hypertrophy of body	Regions of the body are thickened. This additional tissue is part of the body and is enclosed by the hypodermis.
Extensions	Small hypertrophied regions of the body.
Humpback	Extensions, protrusions on the dorsal side of the body. The counterpart, extensions on the ventral side of the body, would be scored as "multi vulva" in the section "Vulva". The distinction between a non vulva-like extension versus a vulva-like extension will be made with a high power microscope.
Truncated body	Part of the body is missing.
Withered body	Part of the body is thinned.
Twisted	Twisted body. The rotation along the anterior-posterior body axis can be seen by the twisted gut/gonad tube or because the vulva and the rectum are not orientated in the same (ventral) direction.
Fat	Worm is thicker and darker than wild type.
Pale	Worm is brighter than wild type.
Pale with dark spots	Worm is brighter than wild type and contains dark spots.
Clear	Worm is nearly transparent.
Full of vacuoles	Worm contains more vacuoles than wild type. Vacuoles have a darker or opal appearance and resemble little moon craters.
Fluid-filled	Liquid flows all over the body.

- 47 -

Poured out	Contents of the worm like the gonad is released through the vulva.
Burst	Dead worm with bursted body shape.

Tail defects

Only tail truncated	Blunt body end; whipe is missing.
Tail shape aberrant	Tail or tail whipe is kinked, shortened or thickened.
Knob-like	Tail whipe has knob-like structures.

Cuticle defects

Blistered	Fluid-filled transparent blisters separated by the hypodermis outside on the body. Clearly different from extensions.
Molting defective	More worms are caught in their old skin like the sloughing of a snake.

It is possible to score body shape phenotypes by image acquisition followed by image analysis. The advantage in the automation of the profiling procedure is the quantification of the strength of a phenotype or the presence of the phenotype in a population. A disadvantage is that the procedure for analysing an image for every possible phenotype may be more elaborate than simply scoring by eye. Furthermore, certain details are difficult to access by video analysis e.g., blister versus protrusions.

Table 10: list of scientific body shape phenotypes, together with their corresponding technical definitions, in terms of characteristics which can be comparatively measured relative to wild-type characteristics using automated measuring apparatus.

Scientific phenotype	Technical definition	Technical phenotype
Proportion abnormal		
Short	Body length less than wild type	Short
Long	Body length more than wild type	Long
Thin	Body diameter less than wild type	Thin

- 48 -

Thick	Body diameter more than wild type	Thick
Dumpy		<i>Disappears</i>
Spindle-shaped		<i>Disappears</i>

Head defects

5	Hypertrophied head	Total head volume has increased	Hypertrophied head
	Extensions on head	Head will be subdivided in n trapezes (or n slices). The diameter of different trapezes can be compared pairwise. The deviation of the diameter can also be located to one side	Extensions on head
	Notched head		Extensions only on one side
	Hammer head		Extensions are pairwise
	Dystrophied head	Total head volume has decreased	Dystrophied head
10	Swollen		<i>Disappears</i>
	Rounded	In the tip trapeze the top diameter is increased	Rounded
	Tapering	The diameter of the tip trapezes are decreased	Tapering
	Vacuoles only in head		<i>Disappears</i>
	Only head bent	The head is most of the time in a certain position that can be measured by an average angle between tip and head/body connection	Tip of head is more often in one position
15	Autodecapitation		<i>Disappears</i>

Example 13**Use of GFP in profiling *C. elegans***

20

A lot of features of *C. elegans* as described in Table 1 can be easily monitored, either automatically by image analysis, microtiter plate readers, or visual means, e.g. by normal microscopy or by Nomarski

25 microscopy. Some features of *C. elegans* are more difficult to visualize. For these characteristics transgenic animals expressing a marker gene are very useful. Moreover, even for characteristics that are rather easily to score, the use of a nematode

30 expressing a marker gene, such as GFP, LacZ, or luciferase, enhances the fingerprinting of *C. elegans*.

- 49 -

The *C. elegans* can be a wild type, a mutant, or a strain subjected to a compound or environmental stress, or a combination of those.

5 *C. elegans* mutant *unc-23* has a fingerprint, which comprises "jerky movement", "tend to coil", "bent head" and "egl". Expressing GFP in the muscle cells of the animal could result in identification and scoring
10 of additional characteristics such as "improperly folded muscles", and/or "detached muscles in head region", and/or "no muscles in head region", and/or "defective muscle attachment", and/or "vulva muscle defects" (data not shown).

15 Similarly, *C. elegans* mutant *unc-71* has a fingerprint which comprise "reduced movement", "weak amplitude", "strong kinker", and "slightly egl". When introducing GFP in the neurons of the animals no apparent extra
20 fingerprint features were observed. A closer look at the neurons of this mutant worm revealed at least following extra phenotypes: "fasculation defects", "VD/DC connection defects" (data not shown).

25 GFP-phenotypes are hence very important in allowing phenotypes which are not otherwise visible to be measurable with Nomarski or dissection microscopy. GFP-phenotypes are further important in the pinpointing of defects to certain tissues and cells, and moreover GFP-phenotypes are important in
30 distinguishing between similar defects with different causes.

- 50 -

Claims:

1. A method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

(a) providing a worm having a defect in at least one gene,

(b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,

(c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said defect,

(d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of worms each of which has a different defect, and

(e) collating the phenotypic profiles so obtained into a library of said profiles.

2. A method as claimed in claim 1 wherein in step (c) at least three changed characteristics are scored.

3. A method as claimed in claim 1 or claim 2 wherein in step (c) at least six changed characteristics are scored.

4. A method as claimed in any preceding claim wherein in step (c) at least ten characteristics are scored.

- 51 -

5. A method as claimed in any preceding claim wherein said worm is *Caenorhabditis elegans*.

6. A method as claimed in any preceding claim wherein steps (a) to (c) are carried out in respect of substantially every gene in the worm genome.

7. A method as claimed in any preceding claim which includes the step of manipulating said worm to generate said defect in said at least one gene.

8. A method as claimed in any preceding claim wherein said defect is selected from the absence of expression of said gene, the reduction in expression of said gene, the over-expression of said gene, the expression of a functionally defective protein, the expression of a truncated protein, the misexpression of a protein, the ectopic misexpression of a protein, the expression of a protein of altered stability or the alteration of gene expression as a function of time.

9. A method as claimed in claim 7 or 8 wherein said manipulation is carried out on wild-type *C. elegans* or a selected mutant thereof.

10. A method as claimed in claim 9 wherein said selected mutant harbours multiple mutations.

11. A method as claimed in claim 7 or 8 wherein said manipulation is carried out on *C. elegans* carrying a reporter gene.

12. A method as claimed in claim 11 wherein said reporter gene is LacZ or green fluorescent protein (GFP).

- 52 -

13. A method as claimed in any one of claims 7 to 12 wherein said manipulation is carried out on a transgenic *C. elegans*.

5 14. A method as claimed in claim 13 wherein said transgenic *C. elegans* expresses a human gene.

15 15. A method as claimed in claim 14 wherein said human gene is a known drug target.

10

16. A method as claimed in claim 14 or claim 15 wherein said human gene is one associated with a human disease.

15

17. A method as claimed in claim 14 or 15 wherein said human gene is a candidate human disease gene.

20

18. A method as claimed in any of claims 7 to 17 wherein said manipulation is carried out on only a sub-set of *C. elegans* cells.

25

19. A method as claimed in any preceding claim wherein changed characteristics in said worm carrying said defect compared to a worm that does not carry said defect are identified by light microscopy, differential interference contrast optics, fluorescence microscopy, immunochemical detection or spectrophotometric detection, radiation detection, calorimetric detection, fluorescence detection or luminescence detection.

30

35

20. A method as claimed in any preceding claim wherein changed characteristics in said worm carrying said defect compared to a worm that does not carry said defect are identified by a pH change or a change in electrical potential.

- 53 -

21. A method as claimed in any preceding claim wherein said plurality of changed characteristics are scored in a predetermined order to generate said phenotypic profile.

5

22. A method as claimed in any preceding claim wherein the scoring of said plurality of changed characteristics is repeated at predetermined intervals of time.

10

23. A method as claimed in any preceding claim wherein said phenotypic profiles are stored electronically.

15

24. A method as claimed in any preceding claim wherein at least one of said plurality of characteristics is selected from the list shown in Table 1.

20

25. A method as claimed in any one of the preceding claims wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

25

26. A method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

30

(a) exposing a worm to a compound,

(b) measuring any changes in identifiable characteristics of said worm as a result of exposure to said compound,

35

(c) systematically scoring a plurality of any

- 54 -

said changed characteristics to establish a phenotypic profile associated with said compound,

5 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different compounds and

10 (e) collating the phenotypic profiles so obtained into a library of said profiles.

27. A method as claimed in claim 26 wherein in step (c) at least three changed characteristics are scored.

15 28. A method as claimed in claim 27 wherein in step (c) at last six changed characteristics are scored.

20 29. A method as claimed in claim 28 wherein in step(c) at least ten changed characteristics are scored.

25 30. A method as claimed in any one of claims 26 to 29 wherein said nematode worm is *C. elegans*.

31. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds has a known pharmacological activity.

30 32. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds is one which is known to interact with a particular biochemical pathway.

35 33. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds has no known pharmacological activity or

- 55 -

biochemical interaction.

34. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds is from a combinatorial library.

35. A method as claimed in any one of claims 26 to 34 wherein said worm to which said compound is exposed is wild-type *C. elegans* or a selected mutant thereof.

36. A method as claimed in claim 35 wherein said selected mutant harbours multiple mutations.

37. A method as claimed in any one of claims 26 to 34 wherein said worm to which said compound is exposed is *C. elegans* carrying a reporter gene.

38. A method as claimed in claim 37 wherein said reporter gene is LacZ or GFP.

39. A method as claimed in any one of claims 26 to 38 wherein said worm to which said compound is exposed is a transgenic *C. elegans*.

40. A method as claimed in claim 39 wherein said transgenic *C. elegans* expresses a human gene.

41. A method as claimed in claim 40 wherein said human gene is a known drug target.

42. A method as claimed in claim 40 wherein said human gene is one associated with a human disease.

43. A method as claimed in claim 40 wherein said human gene is a candidate disease gene.

- 56 -

44. A method as claimed in any one of claims 30 to 43 wherein said worm is exposed to said compound by feeding the worm on bacteria which have been exposed to said compound.

5

45. A method as claimed in claim 44 wherein said bacteria are *E. coli*.

10

46. A method as claimed in any one of claims 26 to 45 wherein said compound is linked to another compound or carrier substance.

15

47. A method as claimed in anyone of claims 26 to 46 wherein any changed characteristics in said worm resulting from exposure to said compound are identified by light microscopy, differential interference contrast optics, fluorescence microscopy, immunochemical detection, spectrophotometric detection, radiation detection, colorimetric detection, fluorescence detection or luminescence detection.

20

25

48. A method as claimed in any one of claims 26 to 47 wherein any changed characteristics in said worm resulting from said compound are identified by a pH change or a change in electrical potential.

30

49. A method as claimed in any one of claims 26 to 48 wherein said plurality of changed characteristics are scored in a predetermined order to generate said profile.

35

50. A method as claimed in any one of claims 26 to 49 wherein the scoring said plurality of changed characteristics is repeated at predetermined time intervals.

- 57 -

51. A method as claimed in any one of claims 26 to 50 wherein said scoring of changed characteristics is carried out using essentially the same scoring protocol as used in a method in accordance with any one of claims 1 to 25.

52. A method as claimed in any one of claims 26 to 51 which comprises the further step of storing the said phenotypic profiles electronically.

53. A method as claimed in any one of claims 26 to 52 wherein at least one of said plurality of characteristics is selected from the list shown in Table 1.

54. A method as claimed in any one of claims 26 to 53 wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

55. A method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

(a) exposing a worm to an environmental change,

(b) measuring any changes in identifiable characteristics as a result of said environmental change,

(c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said change,

- 58 -

(d) simultaneously or sequentially repeating steps (a) to (c) for each of a plurality of different environmental changes and (e) collating the phenotypic profiles so obtained into a library of said profiles.

5

56. A method as claimed in claim 55 wherein in step (c) at least three changed characteristics are scored.

10

57. A method as claimed in claim 56 wherein in step (c) at least six changed characteristics are scored.

15

58. A method as claimed in claim 57 wherein in step (c) at least ten changed characteristics are scored.

20

59. A method as claimed in any of claims 55 to 58 wherein said environmental change is a change in the pH to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a different pH.

25

60. A method as claimed in any one of claims 55 to 58 wherein said environmental change is a change in the osmolarity to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a different osmolarity.

30

61. A method as claimed in any one of claims 55 to 58 wherein said environmental change is a change in the temperature to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a change in temperature.

35

62. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises

- 59 -

exposure to radiation and in step (d) each of said plurality of environmental changes comprises a different level of radiation.

5 63. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises exposure to a virus and in step (d) each of said plurality of environmental changes comprises exposure to a different virus.

10 64. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises exposure to a bacterium and in step (d) each of said plurality of environmental changes comprises exposure to a different bacterium.

15 65. A method as claimed in any one of claims 55 to 64 wherein said worm is *C. elegans*.

20 66. A method as claimed in any one of claims 55 to 65 including a further feature as defined in any one of claims 5 to 54.

25 67. A method as claimed in any one of claims 55 to 66 wherein said scoring of changed characteristics is carried out using essentially the same scoring protocol as used in a method in accordance with claims 1 to 54.

30 68. A method as claimed in any one of claims 55 to 67 wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx
35 pumping, defecation and fertility.

69. A method of constructing a multiple library

- 60 -

of phenotypic profiles of nematode worms which method comprises carrying out all of the methods of claims 1, 26 and 55.

5 70. A method as claimed in claim 69 wherein step (b) of the method of at least one of claims 1, 26 and 55 comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour,
10 mechanotransduction, pharynx pumping, defecation and fertility.

 71. A method of determining the mode of action of a compound which method comprises the steps of;
15

(a) exposing a nematode worm to said compound

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure
20 to said compound,

(c) systematically scoring a plurality of changed characteristics to establish a phenotypic profile associated with said compound and
25

(d) comparing said phenotypic profile with a library of reference phenotypic profiles wherein said library of reference profiles is obtainable by carrying a method in accordance with any of claims 1
30 to 70.

 72. A method of determining whether a compound or combination of compounds interacts with a particular gene or biochemical pathway which method
35 comprises the steps of;

(a) exposing a nematode worm to said compound or

- 61 -

combination of compounds

(b) measuring any changes in identifiable characteristics of said worm as a result of said exposure,

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile associated with said compound or combination of compounds, and

(d) comparing said profile with a library of reference profiles said library of reference profiles being obtainable by carrying out the method of any one of claims 1 to 70.

73. A method of finding an alternative treatment for a human disease which method comprises the steps of:

(a) exposing a nematode worm to a candidate compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

(d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 31.

74. A method of finding a biochemical pathway in

- 62 -

which a compound known to have pharmacological activity acts which method comprises the steps of:

5 (a) exposing a nematode worm to the known compound,

10 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound, and

15 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 32.

20 75. A method of finding a potential new medicinal indication for a compound of known pharmaceutical activity which method comprises the steps of:

25 (a) exposing a nematode worm to the known compound,

30 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

35 (d) comparing said profile with a library of reference profiles, said library of reference profiles

- 63 -

being obtainable by carrying out a method in accordance with any one of claims 1 to 70.

5 76. A method as claimed in claim 75 wherein said library of reference profiles is obtainable by carrying out a method in accordance with any one of claims 24 to 26.

10 77. A method of identifying the mechanism of action of any side effects associated with a compound of known pharmaceutical activity which method comprises the steps of;

15 (a) exposing a nematode worm to the known compound,

20 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

25 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 32 and/or any of claims 1 to 25.

30 78. A method of attributing a particular gene to a particular biochemical pathway in *C. elegans* which method comprises the steps of:

35 (a) exposing a nematode worm to a compound known to operate in a particular biochemical pathway,

 (b) measuring any changes in the identifiable

- 64 -

characteristics of said worm as a result of exposure to said compound

5 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound, and

10 (d) comparing said, profile with a library of reference phenotypic profiles said library of reference profiles being obtainable by carrying out a method in accordance with any one of claims 1 to 25.

15 79. A method as claimed in any of claims 71 to 78 wherein said nematode worm is selected from wild-type *C. elegans*, a mutant *C. elegans* comprising one or more mutations, a *C. elegans* carrying a reporter gene or a transgenic *C. elegans*.

20 80. A method as claimed in claim 79 wherein said transgenic *C. elegans* expresses a human gene.

25 81. A method as claimed in any one of claims 71 to 80 wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

30 82. A method for elucidating biochemical pathways in a nematode worm which method comprises the steps of:

35 (a) generating a defect in at least one gene in said worm,

(b) measuring any changes in identifiable characteristics of said worm compared to a worm

- 65 -

without said defect,

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said defect, and

(d) comparing said profile with a library of reference phenotypic profiles, said library of references profiles being obtainable by carrying out a method in accordance with any one of claims 1 to 25.

83. A method as claimed in claim 82 wherein said nematode worm is selected from wild-type *C. elegans*, a mutant *C. elegans* comprising one or more mutations, a *C. elegans* carrying a reporter gene or a transgenic *C. elegans*.

84. A method as claimed in claim 82 wherein said defect is selected from the absence of expression of said gene, the reduction in expression of said gene, the expression of a functionally defective protein, the expression of a truncated protein, the misexpression of a protein, the ectopic misexpression of a protein, the expression of a protein of altered stability or the alteration of gene expression as a function of time.

85. A method as claimed in any one of claims 82 to 84 wherein at least three, preferably at least six and more preferably at least ten changed characteristics are scored.

86. A method as claimed in any of claims 82 to 85 which includes the features described in any one of claims 19 to 25.

87. A method of constructing a library of

- 66 -

nematode worms which method comprises the steps of:

(a) providing a worm having a defect in at least one gene.

5

(b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,

10

(c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said defect,

15

(d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of worms, and

20

(e) producing a library of said worms each identifiable by their phenotypic profiles.

88. A method as claimed in claim 87 wherein said phenotypic profiles are collated into a library.

25

89. A method as claimed in claim 87 and 88 comprising any one of the features described in any one of claims 2 to 25.

30

90. A method of constructing a library of nematode worms which method comprises the steps of:

(a) exposing a worm to a compound,

35

(b) measuring any changes in identifiable characteristics of said worm as a result of exposure to said compound,

- 67 -

(c) systemically scoring a plurality of any said changed characteristics to establish a phenotypic profile associated with said compound,

5 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different compounds, and producing a library of said worms each identifiable by their phenotypic profiles.

10 91. A method as claimed in claim 90 wherein said phenotypic profiles are collated into a library.

15 92. A method as claimed in claim 90 or 91 comprising any one of the features disclosed in any one of claims 27 to 54.

93. A method of constructing a library of nematode worms which method comprises the steps of:

20 (a) exposing a worm to an environmental change,

(b) measuring any changes in identifiable characteristics as a result of said environmental change,

25 (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said change,

30 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different environmental changes, and

35 (e) producing a library of said worms each identifiable by their phenotypic profile.

94. A method as claimed in claim 93 wherein said

- 68 -

phenotypic profiles are collated into a library.

95. A method as claimed in claim 93 or claim 94 comprising any one of the features disclosed in any one of claims 56 to 70.

96. A method of determining the mode of action of a compound which method comprises the step of:

(a) exposing a nematode worm to said compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile associated with said compounds, and

(d) comparing said phenotypic profile with the library of phenotypic profiles obtainable by the method of any one of claims 88, 91 or 94.

97. A method of determining whether a compound or a combination of compounds interacts with a particular gene or biochemical pathway which method comprises the steps of:

(a) exposing an nematode worm to said compound or combination of compounds,

(b) measuring any changes in identifiable characteristics of said worm as a result of said exposure,

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic

- 69 -

profile associated with said compounds or combination of compounds, and

5 (d) comparing said phenotypic profile with a library of reference profiles wherein said library of reference profiles is obtainable by the method of any one of claims 88, 91 or 94.

10 98. A method of finding an alternative treatment for a human disease which method comprises the steps of:

(a) exposing an nematode worm to a candidate compound,

15 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

20 (c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

25 (d) comparing said profile with a library of 35referenced profiles, wherein said library of referenced profiles is obtainable by carrying out the method in accordance with any one of claims 88, 91 or 94.

30 99. A method of finding a biochemical pathway in which a compound known to have pharmacological activity acts which method comprises the steps of:

35 (a) exposing a nematode worm to the known compound, measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

- 70 -

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

5 (c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

10 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

15 100. A method of finding a potential new medicinal indication for a compound of known pharmaceutical activity which method comprises the steps of:

20 (a) exposing an nematode worm to the known compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

25 (c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

30 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

35 101. A method of identifying the mechanism of action of any side effects associated with a compound of known pharmaceutical activity which method

- 71 -

comprises the steps of:

(a) exposing a nematode worm to the known compound,

5

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

10

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

15

(d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

20

102. A method of attributing a particular gene to a particular biochemical pathway in *C. elegans* which method comprises the steps of:

(a) exposing a nematode worm to a compound known to operate in a particular biochemical pathway,

25

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

30

(c) systemically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

35

(d) comparing said profile with a library of reference phenotypic profiles, said library of reference profiles being obtainable by carrying out the method in accordance with any one of claims 88, 91

- 72 -

or 94.

103. A method as claimed in any one of claims 96 to 102 wherein said nematode worm is selected from wild-type *C. elegans*, a mutant *C. elegans* comprising one or more mutations, a *C. elegans* carrying a reporter gene or a transgenic *C. elegans*.

104. A method as claimed in claim 103 wherein said transgenic *C. elegans* expresses a human gene.

105. A method of establishing a phenotypic profile for a nematode worm which method comprises measuring and scoring at least three, preferably at least six and more preferably at least ten characteristics of said worm which are not exhibited by wild-type worms.

106. A method as claimed in claim 105 wherein said characteristics not exhibited by wild-type worms are selected from the list shown in Table 1.

107. A method as claimed in claim 105 or claim 106 which comprises measuring and scoring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

108. A method as claimed in any one of claims 105 to 107 wherein said phenotypic profile is established for a nematode worm which is selected from a worm having one or more mutations, a worm which has been exposed to a compound or combination of compounds, a transgenic worm, a worm carrying a reporter gene or a worm which has been exposed to an environmental change.

- 73 -

109. A method as claimed in claim 108 wherein said transgenic worm comprises a human gene.

5 110. A method as claimed in claim 108 wherein said compound has known pharmacological activity.

10 111. A method as claimed in claim 108 wherein said compound is known to be active in a particular biochemical pathway.

112. A method as claimed in claim 108 wherein said compound or combination of compounds is from a combinatorial library of compounds.

15 113. A compound which has potential therapeutic activity in a mammal which has been identified in a method as claimed in any one of claims 71 to 81 or 96 to 104.

20 114. A library of nematode worms obtainable by a method as claimed in any one of claims 87 to 95.

25 115. A library as claimed in claim 114 wherein said nematode worm is *C. elegans*.

FIG. 1.





FIG. 2



PCTLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 1/04, 1/00, 15/01, 15/10	A3	(11) International Publication Number: WO 00/34438 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/EP99/09710 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 9826890.7 7 December 1998 (07.12.98) GB (71) Applicant (for all designated States except US): DEVGEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KALETTA, Titus [BE/BE]; (BE). FEICHTINGER, Richard [BE/BE]; (BE). VAN POUCKE, Jonas [BE/BE]; (BE). VAN GEEL, Anton [BE/BE]; (BE). APPELMANS, Saskia [BE/BE]; (BE). VAN CRIEKINGE, Wim [BE/BE]; (BE). BOGAERT, Thierry [BE/BE]; Devgen NV, Technologiepark 9, B-9052 Zwijnaarde (BE). (74) Agent: BOULT WADE TENNANT; 27 Fumival Street, London, EC4A 1PQ (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 9 November 2000 (09.11.00)
(54) Title: METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC PROFILES (57) Abstract Methods are provided for use in constructing libraries of phenotypic profiles in a nematode worm such as <i>C. elegans</i> . The methods require measurement of identifiable characteristics of the worm and systematic scoring of these characteristics. Also provided are methods of identifying compounds with potential pharmacological activity, for determining the mode of action of a given compound and for assigning genes to particular biochemical pathways.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/99/09710

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N1/04 C12N1/00 C12N15/01 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 90 09096 A (CAMBRIDGE NEUROSCIENCE RES ;HORVITZ HOWARD ROBERT (US)) 23 August 1990 (1990-08-23) Cited against inventions 1 and 2 in their entirety and inventions 3 and 4 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. page 7, line 18 -page 8, line 23 page 15, line 14 - line 30 --- -/--</p>	<p>1-112, 114,115</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

25.07.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP/97/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KATSURA ET AL.: "Isolation, characterization and epistasis of fluoride-resistant mutants of <i>Caenorhabditis elegans</i> " GENETICS, vol. 136, 1994, pages 145-154, XP000886900 Cited against invention 1 abstract; tables 1-4 page 145, column 1 -page 146, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
X	VAN SWINDEREN ET AL.: "Quantitative trait loci controlling halothane sensitivity in <i>Caenorhabditis elegans</i> " PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 8232-8237, XP002137784 Cited against invention 2 in its entirety and invention 3 insofar as "environmental changes" can also include those changes due to (e.g. toxic) compounds. abstract page 8232, column 1 -page 8233, column 1 ---	1-25,71, 72, 75-89, 96-112, 114,115
A	AHRINGER ET AL.: "Turn to the worm!" CURRENT OPINION IN GENETICS AND DEVELOPMENT, vol. 7, 1997, pages 410-415, XP000886904 cited in the application Cited for all inventions the whole document ---	1-112, 114,115
X	WO 96 38555 A (BOGAERT THIERRY ;STRINGHAM EVE (CA); VANDEKERCKHOVE JOEL (BE)) 5 December 1996 (1996-12-05) Cited against inventions 2 and 3 page 35, line 22 -page 36, line 28; claim 43 ---	26-68, 71-77, 79-81, 90-114
A	SAMOILOFF, M.R. ET AL: "The use of nematodes in marine ecotoxicology. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 1." MAR. TOX., (1984) PP. 407-426. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000812-8., XP000886947 Dep. Zool., Univ. Manitoba, Winnipeg, Man. R3T 2N2, Canada Cited for inventions 3 and 4 page 413, paragraph 2 ---	55-68, 71,72, 75, 79-81, 93-112, 114,115

-/--

INTERNATIONAL SEARCH REPORT

international application no
PCT/EP/09710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BOGAERT, T. ET AL: "Determination of the toxicity of four heavy metal compounds and three carcinogens using two marine nematode species, Monhystera microphthalma and Diplolaimelloides brucieii. ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. VOL. 2." MAR. TOX., (1984) PP. 21-30. MEETING INFO.: INTERNATIONAL SYMPOSIUM ON ECOTOXICOLOGICAL TESTING FOR THE MARINE ENVIRONMENT. GHENT (BELGIUM). 12-14 SEP 1983. ISSN: 90-9000814-4; 90-9000813-6., XP000886948</p> <p>Lab. Mol. Biol., Med. Res. Counc. Cent., University Med. Sch., Hills Rd., Cambridge CB2 2QH, UK</p> <p>Cited for inventions 3 and 4 the whole document</p> <p>-----</p>	<p>55-68, 71,72, 75, 79-81, 93-112, 114,115</p>

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 99/09710

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 113
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-25,78,82-89 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound or gene, comprising comparing the phenotypic response of a nematode treated with said compound or with a defect in said gene with a library of multiple phenotypic traits of nematodes with genetic defects and subject-matter relating thereto.

2. Claims: 26-54,73,74,90-92 completely; 71,72,75-77,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes treated with other compounds and subject-matter relating thereto.

3. Claims: 55-68,93-95 completely; 71,72,75,79-81, 96-115 partially

Method for determining the mode of action of a compound, comprising comparing the phenotypic response of a nematode treated with the compound with a library of multiple phenotypic responses of nematodes subjected to environmental changes and subject-matter relating thereto.

4. Claims: 69,70 completely; 71,72,75,79-81,96-113 partially

Method for determining the mode of action of a compound or gene, comprising the methods of inventions 1-3 referred to above and subject-matter relating thereto.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box 1.2

Claims Nos.: 113

It is not possible to carry out a meaningful search into the state of the art on the basis of claim 113 because its subject-matter ("agonists" and "antagonists") is structurally undefined and could not in any event have been functionally tested in the prior art.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/EP 99/09710

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9009096	A	23-08-1990	AU	5106790 A	05-09-1990

WO 9638555	A	05-12-1996	AU	6123496 A	18-12-1996
			EP	0832222 A	01-04-1998

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 1/04, 1/00, 15/01, 15/10	A2	(11) International Publication Number: WO 00/34438 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/EP99/09710 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 9826890.7 7 December 1998 (07.12.98) GB (71) Applicant (for all designated States except US): DEVGEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): KALETTA, Titus [BE/BE]; (BE). FEICHTINGER, Richard [BE/BE]; (BE). VAN POUCKE, Jonas [BE/BE]; (BE). VAN GEEL, Anton [BE/BE]; (BE). APPELMANS, Saskia [BE/BE]; (BE). VAN CRIEKINGE, Wim [BE/BE]; (BE). BOGAERT, Thierry [BE/BE]; Devgen NV, Technologiepark 9, B-9052 Zwijnaarde (BE). (74) Agent: BOULT WADE TENNANT; 27 Furnival Street, London, EC4A 3DF (GB). <div style="text-align: center;">05 JUN 2001</div>		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC PROFILES (57) Abstract Methods are provided for use in constructing libraries of phenotypic profiles in a nematode worm such as <i>C. elegans</i> . The methods require measurement of identifiable characteristics of the worm and systematic scoring of these characteristics. Also provided are methods of identifying compounds with potential pharmacological activity, for determining the mode of action of a given compound and for assigning genes to particular biochemical pathways.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD FOR CONSTRUCTING LIBRARIES OF PHENOTYPIC
PROFILES

5 The present invention is concerned with the field
of 'genetic pharmacology'. Specifically, it relates to
methods which can determine, among other things,
whether a compound has potential pharmacological
activity, whether a compound interacts with a
particular gene or biochemical pathway in man or
10 animals, what side effects are likely to be associated
with a particular pharmaceutical compound and/or the
mode or modes of action of any compound with
biological activity. Additional uses for the methods
of the invention include the assignment of function to
15 particular genes or assignment of genes and their
encoded proteins to particular biochemical pathways.
In particular, the invention relates to the use of a
microscopic nematode worm, for example *Caenorhabditis*
elegans, and libraries of such worms in the
20 aforementioned methods. These new methods are able to
enhance and accelerate the drug discovery process.

Prior to the early 1990's the search for new
compounds having the potential to combat human or
animal disease was often begun by taking a compound
25 known to have a particular pharmacological activity,
synthesising structurally related variants and then
testing those variants against the known target.

The test against the target might be carried out
in vivo, for example by use of animal models of a
30 human disease. Alternatively, if a particular molecule
was known to be implicated in the progress of a
disease, the compounds could be tested for
interaction with the molecule *in vitro*. The
limitations of such methods are that in the event of a
35 negative result no other information about the
pharmaceutical potential of the compound tested is

- 2 -

gained. For example, an *in vitro* test might show a compound to have no inhibitory action against a particular target enzyme but that compound might have an inhibitory action against another enzyme in the same biochemical pathway as the target enzyme and therefore, in fact, have potential in treatment of the target disease. Animal tests, while providing a reasonable indication of both efficacy and toxicity, provide no information at all about the mode of action of the compound, and therefore the possible reasons for any toxicity. Furthermore, they are time-consuming and expensive and do not lend themselves to automation.

Since the early nineties there have been two developments in particular which have revolutionized the drug discovery process, these being the new sciences of 'genomics' and 'combinatorial chemistry'. It has now been realised that a vast number of diseases have a genetic component and they are not purely the result of environmental influences. Indeed, it is possible that nearly all diseases are multifactorial and will have some degree of genetic basis, albeit very small in some cases. A huge amount of effort is being directed at the present time to the study of the organisation of the genomes of various unicellular and multicellular organisms, including humans. This involves the identification and sequencing of all the genes in a particular genome. Such activity does not only allow for hunting of genes which are directly associated with particular diseases but each of the genes found and the proteins they encode can become, directly or indirectly, a target against which compounds can be screened, whether or not that gene has yet been associated with a disease or indeed has any identified function at all.

- 3 -

Furthermore, rather than starting from a compound of known 'activity' and relying on theoretical structure/function relationships to synthesise new candidate compounds, vast libraries of compounds, of uniform activity can be very rapidly synthesized in an automated manner by combinatorial chemistry. Thus, there is now potential to screen thousands of compounds against thousands of genes and the proteins they encode in very rapid high throughput screens (HTS) and to link compounds to genes and genes to disease.

The present inventors have discovered that these new technologies for drug discovery can conveniently be married with a particular multicellular organism, a nematode worm, *C.elegans*, which has been well characterised genetically and morphologically. They have thereby developed new methods, which are extremely powerful, rapid and convenient and can play an essential part in a drug discovery program.

C. elegans is a microscopic nematode worm which occurs naturally in the soil but can be easily grown in the laboratory on nutrient agar inoculated with bacteria, preferably *E. coli*, on which it feeds. Each worm grows from an embryo to an adult worm of about 1 mm long in three days or so. As it is fully transparent at all stages of its life, cell divisions, migrations and differentiation can be seen in live animals. Furthermore, although its anatomy is simple its somatic cells represent most major differentiated tissue type including muscles, neurons, intestine and epidermis. Accordingly, differences in phenotype which represent a departure from that of a wild-type worm are relatively easily observed, either directly by microscopy or by using selective staining procedures, and many of these phenotypic differences submit to quantitative measurement. Many *C. elegans* mutants have

- 4 -

been identified and their phenotypes described, for example, see *C. elegans* II Ed. Riddle, Blumenthal, Meyer and Priess, Cold Spring Harbor Laboratory Press, 1997. The *C. elegans* genome is now almost entirely
5 sequenced as a result of the *C. elegans* genome project, carried out at the Sanger Center and Washington University School of Medicine. The sequence is available in a public database at
http://www.sanger.ac.uk/projects/C_elegans/. As a
10 result of this it has emerged that *C. elegans* comprises genes which have equivalents that are widely distributed in most or all animals including humans.

Methods for creating mutant worms with mutations in selected *C. elegans* genes are known in the art, for
15 example see J. Sutton and J. Hodgkin in 'The Nematode *Caenorhabditis elegans*' Ed. By William B. Wood and the Community of *C. elegans* Researchers CSHL, 1988 594-595; Zwaal et al; Target-Selected Gene Inactivation in *Caenorhabditis elegans* by using a Frozen Transposon
20 Insertion Mutant Bank' 1993, Proc. Natl. Acad. Sci. USA 90 pp 7431-7435; Fire et al, Potent and Specific Genetic Interference by Double-Stranded RNA in *Caenorhabditis elegans* 1998, Nature 391 860-811.

The possibility that *C. elegans* might be useful
25 for establishing links between compounds and specific *C. elegans* genes by virtue of comparison of phenotypes generated by exposure to particular compounds and by selected mutations is considered by Rand and Johnson in Methods of Cell Biology, Chapter 8, vol 84,
30 *Caenorhabditis elegans*: Modern Biological Analysis of an Organism Ed. Epstein and Shakes, Academic Press, 1995 and J. Ahringer in Curr. Op. in Gen. & Dev. 7; 1997; 410-415.

However, these authors observe and attribute
35 altered phenotypes on the basis of a single changed characteristic such as, for example, pharyngeal

- 5 -

pumping rate or defecation frequency. Since that single characteristic may be determined by expression of a number of genes and the operation of several biochemical pathways such a crude assessment of phenotype is not sufficient to establish a link between any one gene or pathway and a compound to which the worm has been exposed. As such the procedure would not be sensitive enough for resolution of the properties of thousands of compounds in a high throughput compound screen. An additional problem with the proposals of the prior art is that known phenotypic characteristics have all been described differently by different workers in the *C. elegans* field. Phenotype descriptions in the literature largely omit aspects not directly related to or not recognised to be related to the principle interest of the individual researcher. There is no standard nomenclature to identify a specific change. Without this it is impossible to equate newly observed phenotypes with particular known phenotypes for comparison purposes.

The present inventors have developed methods which solve these problems and thereby have converted *C. elegans* into a really useful tool in the drug discovery field. Specifically, in respect of each worm a 'phenotype profile' or 'fingerprint' is established based on looking for plurality of changed characteristics in a particular mutant or worm which has been exposed to an environmental change or a compound. Furthermore, each profile is scored by following a strict standard protocol of measurement and a standard description is applied to each characteristic. The determination of a phenotypic profile in this way for a plurality of mutants or worms exposed to compounds illuminates differences between different mutants or otherwise treated worms

- 6 -

which would not be apparent based on prior art methods. Furthermore, the standard scoring protocol and nomenclature allows the phenotypic profiles obtained to be collated into a library of reference profiles for direct comparison purposes. Thus, libraries of reference profiles can be established for mutant worms and for worms exposed to particular environmental changes or different sorts of compounds. Such libraries allow complex patterns of linkage to be established between particular compounds and particular genes or biochemical pathways and between individual compounds of known or unknown biochemical or pharmacological activity.

In accordance with a first aspect of the present invention there is provided a method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

- (a) providing a worm having a defect in at last one gene.
- (b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,
- (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotype profile associated with said defect,
- (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of worms each of which has a different defect, and
- (e) collating the phenotypic profiles so obtained into a library of said profiles.

- 7 -

Caenorhabditis elegans is the preferred nematode worm although the method could be carried out with other nematodes and in particular with other microscopic nematodes, preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size are extremely suited for use in mid- to high-throughput screening as they can easily be grown in the wells of a multi-well plate of the type generally used in the art to perform such screening.

It is preferred to establish the phenotypic profile on the basis of the measurement and scoring of at least three different characteristics, preferably at least six characteristics and more preferably at least ten characteristics. It will be appreciated that the more differences which can be scored between a worm with a genetic defect and a worm without the defect the better the resolution between different mutants. Although not limited to such, at least one of the plurality of changed characteristics which can be measured and scored may be selected from the list shown in Table 1, and possibly each of all the changed characteristics scored is one of those shown in Table 1.

In a preferred embodiment, the method used to establish the phenotypic profile comprises measurement and scoring of two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility. This list provides a core set of measurable characteristics which can be used to establish an informative phenotypic profile for any type of worm. Furthermore,

- 8 -

each of these characteristics is measurable using technical measuring apparatus, such as video image analysis, multiwell plate reader, and/or a technical assay procedure. In the most preferred embodiment, the method used to establish the phenotypic profile comprises measurement and scoring of all eight of the listed core characteristics. Measuring and scoring this set of core characteristics allows meaningful comparisons to be made between phenotypic profiles for worms subjected to diverse interventions. AS exemplified herein, comparisons can be drawn between profiles for two different mutant worms and between profiles for mutant worms and profiles for worms exposed to compound.

It is to be understood the terms "measuring" or "measurement" as used in connection with any of the methods described and claimed herein are to be interpreted as including not just absolute quantitative measurement wherein a numerical value is assigned to the characteristic but also comparative measurement, wherein characteristics of a worm which has been subject to an intervention (i.e. mutation, exposure to compound, exposure to environmental change) are measured relative to the same characteristics of a wild-type worm and scored as being 'larger', 'smaller', 'longer', 'shorter', 'fatter', 'thinner', 'darker', 'paler' etc.

For comparison purposes it is essential that the scored characteristics are represented in the same order for each profile. For standardization of procedure between different workers or to facilitate automation, measurement and scoring of the characteristics could be carried out in a pre-determined order according to a standard protocol. However, this is not essential to the operation of the method. In its simplest form and as shown in Example 5, the characteristics are recorded in a binary manner

- 9 -

as 'present' or 'not present' based on deviations from wild-type worms.

It is desirable to establish a library which comprises a phenotypic profile in respect of a defect in each gene in the worm genome and/or different defects in the same gene (allelic variations). As
5 aforesaid there are a considerable number of available mutants (see Riddle, Blumenthal, Meyer and Priess and Ahringer above). In addition new ones can be generated
10 by specific gene and site directed mutation and knock-out methods known to those skilled in the art such as ethyl methanesulphonate (EMS) mutagenesis, transposon insertion or genetic interference using double
stranded RNA (see Sutton and Hodgkin, Zwaal et al and
15 Fire et al above). The known or newly generated genetic defects may manifest themselves, for example, as the absence of expression of a gene, the reduction in expression of a gene, the over-expression of a
gene, the expression of a functionally defective
20 protein, the mis-expression of a protein, the ectopic mis-expression of a protein, the expression of a protein of altered stability or the alteration of gene expression as a function of time.

Generally, the manipulation of *C. elegans* to
25 generate genetic defects can be carried out on wild-type worms or worms with existing single or multiple mutations. It may be desirable to genetically manipulate *C. elegans* carrying a reporter gene construct. The reporter molecule might be LacZ or
30 green fluorescent protein but many other reporter molecules are known to those skilled in the art. Reporter gene constructs for *C. elegans* are described in Chalfie et al, 1994, Science 263 pp 802-805. It can
also be desirable to genetically manipulate and then
35 profile a transgenic worm, preferably a worm carrying a human gene, particularly where the gene is

- 10 -

associated with, or is a candidate for association with a human disease and therefore a putative drug target. A list of human diseases for which a particular gene has been implicated is given in the paper by J. Ahringer (see above) and also provided by OMIM. Center for Medical Genetics, John Hopkins University and National Biotechnology Information, National Library of Medicine, 1996. <http://www.ncbi.nlm.nih.gov/omim/>, although these lists are not necessarily exhaustive.

It is easy to establish transgenic lines in *C. elegans* and the methodology is described in Craig Mello and Andrew Fire, *Methods in Cell Biology*, Vol 48 Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

A form of the worm which may show a change in phenotype and may therefore be subject to profiling as described above is one in which the genetic defect and/or transgene and/or reporter gene is only present in a sub-set of the cells of the worm. It is possible for just the cells of a particular tissue to be the subject of a genetic manipulation.

The worm which is to be subject to determination of its phenotypic profile can be cultured by methods well-known in the art. *C. elegans* can grow on nutrient agar which has first been inoculated with bacteria on which the worms feed. Suitable culture methods are described in Rand and Johnson (see above) and in the examples given herein. Measurement of any changed characteristics which will determine the profile may be carried out using light microscopy, differential interference contrast optics or fluorescence microscopy. In addition immuno-chemical detection, colorimetric detection or detection of fluorescence, luminescence or radioactive labels may be used. In

- 11 -

some cases the changed characteristics may be biochemical only and might be detected, for example by a pH change in the growth media or a change in electrical potential. Different characteristics may need to be determined at different points in the growth cycle of the worm. For example, some phenotypic characteristics may be manifested only in the larvae while others are only detectable in the adult worm. In some cases it may be necessary to make several measurements of the same characteristic at pre-determined time intervals.

Phenotypic profiles generated by the methods described above can be collated into a library of profiles which are stored electronically on a database. However, it will be appreciated that the invention also provides a method of constructing a physical library or bank of worms each identifiable by their individual phenotypic profile. Such a worm library can be created using any or all of the methods described above and used for comparative purposes. The worms may be maintained by the culture methods described herein and/or frozen for long term storage by methods known to those skilled in the art. Libraries of phenotypic profiles or fingerprints of mutant worms or mutant worm libraries can be used to determine linkages between different genes and hence identify biochemical pathways. A particularly important use is the profiling of several mutations of the same gene and several genes of the same pathway. Different mutations in the same gene can have different phenotypes and often it is found that a careful analysis of the allelic series of a gene reveals important information that is hidden under a more severe phenotype of a null mutant (complete knock out, e.g. if it is lethal). Phenotypic profiles of different mutations of the same gene allow

- 12 -

characterisation of the gene by simply combining
(logical OR) the profiles of all the mutations,
whether they have been generated at the same time or
not. It is possible, however, to handle the mutations
5 separately and make more detailed connections, for
example, concerning protein domains in case the
similarity of phenotypes cluster with the sites of the
mutations.

Described above are methods for constructing a
10 library of phenotypic profiles for worms with a
plurality of genetic defects or a library of mutant
worms. However, in accordance with a second aspect the
present invention provides a method of constructing a
library of phenotypic profiles of nematode worms which
15 comprises the steps of:

- (a) exposing a worm to a compound,
- (b) measuring any changes in identifiable
20 characteristics of said worm as a result of
exposure to said compound,
- (c) systematically scoring a plurality of any
25 said changed characteristics to establish a
phenotypic profile associated with said
compound,
- (d) simultaneously or sequentially repeating
30 steps (a) to (c) in respect of each of a
plurality of different compounds, and
- (e) collating the phenotypic profiles so
obtained into a library of said profiles.

35 Methods for culturing *C. elegans* in the presence
of a test compound are described by Rand and Johnson

- 13 -

mentioned above and in the examples herein. In its simplest form a solution of the compound in a suitable solvent may be spread over a bacterial lawn on an agar plate before inoculation with the worm. Additional
5 refinements include feeding the worm with bacteria, preferably *E. coli*, which have taken up the compound or attaching the compound to a carrier compound which is particularly attractive to the worm.

The worms which are exposed to the compound may
10 be wild-type worms, mutant worms, transgenic worms and/or worms carrying reporter gene constructs as already described herein. Further the measurement and scoring of a plurality of changed characteristics is carried out by exactly the same procedures as already
15 described herein for the phenotypic profiling of mutant worms. This must be a standard format in order that direct comparisons can be made between profiles obtained on exposure to compounds and profiles exhibited by mutants.

20 With compound screening it is possible to build up a series of different libraries depending on the compounds being tested. For example one library can comprise profiles generated in respect of each of the known compounds in a Pharmacopoeia, in other words
25 compounds with known pharmacological activity.

Another library can comprise profiles generated by compounds known to interact with a particular biochemical pathway, which may or may overlap with those compounds from the Pharmacopoeia. Other
30 libraries could include profiles for known compounds but with no known biological activity or compounds which are completely new molecules such as might be generated by combinatorial chemistry. As aforesaid the present invention is not limited to the production
35 of phenotypic profile libraries but includes libraries or banks of worms whose phenotypic profile has been altered by exposure to compounds. In particular

- 14 -

embodiments assays may be carried out with several concentrations of the same compound, and/or with mixtures of compounds. For example compounds from compound libraries may each be tested individually or with one or more other influencing compounds. Furthermore, such compound testing protocols may be executed against identical worms or multiple mutant and/or transgenic backgrounds. In a particular example a panel of worm strains, covering a wide range of biochemical pathways and cellular activities by means of mutations in particular pathways, as well as reporter genes, is used for testing compounds. For each compound, potentially at several concentrations, a profile is recorded for the measurable phenotypes of each of the worm strains, either in parallel or sequentially.

In a third of its aspects the invention provides a method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

- (a) exposing a worm to an environmental change,
- (b) measuring any changes in identifiable characteristics as a result of said environmental change,
- (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said change,
- (d) simultaneously or sequentially repeating steps (a) to (c) for each of a plurality of different environmental changes, and

- 15 -

- (e) collating the phenotype profiles so obtained into a library of said profiles.

5 The environmental change may be, for example, a change in pH, osmolarity, temperature, exposure to radiation or exposure to bacteria or viruses. Each of these external influences may result in the manifestation of a different phenotypic profile of characteristics so that libraries of said profiles and affected worms can be constructed. Again, measurements and scoring of the profile should follow a standard protocol in order that valid comparisons can be made between these profiles and those in mutant and compound libraries.

15 The construction of worm and phenotypic profile libraries by the methods described above using the novel phenotypic profiling method described herein provides a very powerful tool for the discovery of new drugs. Profiles in each of the different libraries can be compared and links established between *C. elegans* genes and pathways, compounds and environmental effects. Preferably, the process of measuring and scoring the changed characteristics which go to make up the phenotypic profile is automated, making use of technical measuring apparatus. The profiles so generated may advantageously be stored electronically. Libraries of profiles can then be searched by computer which can identify identical or similar profiles, either within or between the different libraries. Quantitative data calculations, optionally in combination with boolean operations can be used.

30 A comparison of the profile generated by a particular compound with the profiles of particular mutants may indicate the likely gene or biochemical pathway with which the compound interacts in the worm. Other databases can then be searched for a match of

- 16 -

the worm gene with an equivalent human gene. The human gene might already be associated with a human disease as could be determined for example, from the OMIM database mentioned above. Thus, by use of the worm screen a potential candidate drug can be identified. The discovery of the mode of action of a compound with known pharmacological or biochemical activity is facilitated by comparing its phenotypic profile in the worm with the mutant library or environmental change library of profiles to identify possible targets for the compound. other possibilities include finding a new potential medical indication of a known compound, a medical indication for a novel compound, an alternative method of treatment of a known disease or an indication of the reason for the side effect exhibited by some known pharmaceuticals. Testing worms with compounds, scoring the phenotypic profile in the novel manner described herein and then searching previously established libraries of profiles can potentially achieve all those goals. Once a compound has been identified as having the potential to be a therapeutic agent it can be processed through the more traditional drug discovery routes. The compound can be tested in more specific in vitro tests based on the new knowledge of the target for the compound and in animal models of the target disease. Structural variants then can be generated by medicinal chemistry with a view to improving activity.

The invention will now be described with reference to the following Examples, together with accompanying Figures, in which:

Figure 1 is a schematic diagram of the left lateral view of the body of *C. elegans*. The body of *C. elegans* is divided into a head, a body and a tail region. The head region stops at the end of the

- 17 -

pharynx, the body stops at the rectum and the tail includes the tail whipe. *C. elegans* usually crawl on the right side. The ventral located vulva defines the ventral side of *C. elegans*.

- 5 Figure 2 is a schematic diagram of *C. elegans* showing the characteristics "hypertrophy of the head and "extensions on head".

Example 1

10 General Profiling by Plate Drop Assay

- 4ml NGM agar (see 'The Nematode *Caenorhabditis Elegans*' Ed. by William B. Wood and the Community of *C. elegans* Researchers CSHL, 1988, pg 589) is poured
15 into 3cm plate, and seeded with approximately 5µl of an *E. coli* overnight culture and grown preferably for one week at room temperature. If a compound is to be profiled 10µl of compound dissolved in DMSO or other appropriate solution is pipetted onto the bacterial
20 lawn. The lawn should be covered completely. (This step can be omitted if a mutant, transgenic or other worm is being profiled without compound). After overnight soaking in of compound one *C. elegans* (L4 stage) per plate is put in the bacterial lawn. Worms
25 are checked after some hours, plates are incubated at 21°C and worms screened for phenotypes (control have L1 progeny growing). Plates are checked again after 4 days for phenotypes of F1 progeny (control shows all stages up to gravid hermaphrodites). Plates which have
30 to be looked at again on subsequent days because of slow growth or for further checks are put aside. A plate protocol sheet such as that shown in Table 2 is completed deciding on one of the following routes: no effect/unspecific effect/needs to be applied at lower
35 concentrations/needs to be profiled. If concentrations are appropriate and a decision can be made scoring of

characteristics to produce a profile can be started using the profiling list in Table 1. Because the compound is pipetted onto a bacterial lawn rather than it being incorporated into the agar, as has been done in the prior art, this method is designated a 'plate drop assay'.

Table 1

1. Compound specific phenotypes

[illegible]

2. Viability

[illegible]

- 19 -

	2.3 Embryonic arrest of F1									
	2.3.1 Leakyness									
	2.3.2 Appearance of eggs									
5	2.3.2.1 dark eggs									
	2.3.2.2 bright eggs									
	2.3.2.3 two-fold or older									
	2.3.2.4 irregular egg size									
	2.4 Larval arrest of F1									
10	2.4.1 Leakyness									
	2.4.2 at L1									
	2.4.3 at L2									
	2.4.4 at L3									
	2.4.5 at L4									
15	2.5 Embryonic arrest of F2									
	2.5.1 Leakyness									
	2.5.2 Appearance of eggs									
	2.5.2.1 irregular egg size									
	2.6 Larval arrest of F2									
20	2.6.1 Leakyness									
	2.6.2 at L1									
	2.7 Died during adulthood (F1)									
	2.8 Died during adulthood (F2)									

25 3. Life cycle

	Phenotype									Comment
	Abnormal									
	3.1 Growth abnormal									
30	3.1.1 only generation cycle slowed down									
	3.1.1.1 oldest stage L1									
	3.1.1.2 oldest stage L2									
	3.1.1.3 oldest stage L3									
35	3.1.1.4 oldest stage L4									
	3.1.2 generation cycle slowed down while displaying defects									
	3.1.2.1 oldest stage L1									
	3.1.2.2 oldest stage L2									
40	3.1.2.3 oldest stage L3									
	3.1.2.4 oldest stage L4									
	3.1.3 stage changed									
	3.1.3.1 delayed hatching									
	3.1.3.2 arrested growth in L1									
	3.2 Dauer formation defective									
45	3.2.1 constitutive dauer									
	3.2.2 non-conditional constitutive									
	3.2.3 defective									
	3.2.4 dies on recovery									
	3.3 Life span changed									
50	3.3.1 Life span is shorter									
	3.3.2 Life span is prolonged									

[illegible]15 **5. Movement**[illegible]

5

6. Mechanotransduction (Touch with a wire and with eyelash)

25

- 23 -

7. Sensory system

Phenotype									Comment
Abnormal									
7.1 Avoidance of bacteria									
7.2 Bordering behaviour									
7.3 Chemotaxis defective									
7.3.1 attraction									
7.3.2 avoidance									
7.4 Thermotaxis defective									
7.4.1 attraction									
7.4.2 avoidance									

8. Environmental response

Phenotype									Comment
Abnormal									
8.1 Osmolarity sensitive									
8.2 Thermotolerance changed									
8.3 UV Resistance changed									
8.4 Oxygen sensitive									

9. Pharynx

Phenotype									Comment
Abnormal									
9.1 Pharynx stuffed									
9.2 Morphology defects									
9.3 Pumping defects									
9.3.1 pumping reduced									
9.3.2 pumping enhanced									
9.3.3 pumping irregular									
9.3.4 no pumping									
9.4 Eating defective									

10. Intestine

Phenotype									Comment
Abnormal									
10.1 Morphology defects									
10.1.1 enlarged									
10.1.2 detached									
10.2 Color of contents									
10.2.1 darker									
10.2.2 lighter									

11. Rectum

[illegible]

20 **12. Gonad**

	Phenotype								Comment
	Abnormal								
	12.1 Morphology defects								
25	12.1.1 defective gonad								
	12.1.2 one arm missing								
	12.1.3 multiple gonad								
	12.1.4 monopolar gonad forward								
	12.1.5 monopolar gonad backward								
30	12.1.6 no gonad								
	12.2 Light brown								

13. Vulva

[illegible]

14. Fertility

[illegible]

- 25 -

5	14.2.4	egg laying defective									
	14.2.4.1	weak Egl									
	14.2.4.2	strong Egl									
	14.2.5	bloated worms									
	14.2.5.1	weak bloating									
	14.2.5.2	strong bloating									
	14.2.5.3	bags of worms									
	14.2.6	no egg laying									
10	14.3	Only oocytes									
	14.4	Sterile									
	14.5	Maternal effect sterile									

15. Male

15	Phenotype										Comment
	Abnormal										
	15.1 Frequency										
	15.1.1	high incidence of males									
	15.2 Mating defective										
20	15.3 Morphology										
	15.3.1	leptoderan tail									
	15.3.2	scrawny									
	15.3.3	copulatory plug									
	15.4 Mating behaviour										
25	15.4.1	defective sensory contact									
	15.4.1.1	no response to dorsal contact									
	15.4.1.2	no response to ventral contact									
	15.4.2	defective backing									
	15.4.2.1	no backing									
	15.4.2.2	no continued backing									
30	15.4.3	defective turning									
	15.4.3.1	loose turns									
	15.4.3.2	stop at the tail									
	15.4.3.3	slide off the tail									
35	15.4.4	defective vulval location									
	15.4.5	defective spicule insertion									

16. Progression of phenotype

40	Phenotype										Comment
	Abnormal										
	16.1 Dependent on generation										
	16.1.1	F1 different from P0									
	16.1.1.1	weaker									
	16.1.1.2	worse									
45	16.1.1.3	lower penetrance									
	16.1.1.4	higher penetrance									
	16.1.1.5	not affected									
	16.1.2	F1 different from F2									
	16.2 Dependent on stage										
50	16.2.1	appearance of phenotype									
	16.2.1.1	after L2									
	16.2.1.2	during adulthood									
	16.2.2	shift of phenotype									
	16.3 Dependent on age										
55	16.3.1	phenotype gets worse									
	16.3.2	phenotype gets better									

Table 2

plate	well	by	date
negative control	positive control	finished	confirmed (≥ 3 worms)
no effect	unspecific effect	needs to be applied at lower concentrations	needs to be profiled

Day 0

compound
invisible
coloured
droplets
crystals
complete crust

bacteria
normal lawn
grown as ring
thin
crust
died

worm
happy
run away
irregular movement
slow movement
no movement

Day 1

appearance
healthy
slightly unhealthy
slightly starved
strong starved
very sick

worm gone
lost
suicide
in agar
starved outside
died in compound

replaced by
number and stage
left progeny

movement
normal
tracks more outside
tracks not in center
amplitude increased loopy
amplitude variable
amplitude decreased
enhanced movement
slow movement
no movement
specific

body
normal gravid adult
pumping defects
light brown messy gonad
pale with dark spots
few eggs in gonad
pharynx stuffed
foregut filled large
hindgut constipated
protruding vulva
other:

progeny
normal
reduced broodsize
younger staged
oocytes
coagulated eggs
dead eggs
dying hatchlings
crippled larvae

Day 4

food
still plenty of
already finished
finished soon
outside comp.
not eatable, died

adult viability
still fertile
laying oocytes
died
died as bag of worms
missing

growth rate
normal
reduced broodsize
younger staged

movement
normal
population more outside
population not in center
amplitude increase, loopy
amplitude variable
amplitude decreased
enhanced movement
slow movement
no movement
specific:

body
normal gravid adult
pumping defects
light brown messy gonad
pale with dark spots
few eggs in gonad
pharynx stuffed
foregut filled large
hindgut constipated
protruding vulva
other:

brood viability
dead eggs
dead larvae
larval arrest
later scoring
day of screen
day of worm

comparison of phenotypes

progeny shows PC phenotype
similar
worse
a few only
weaker
no effect

new worms show phenotype
similar
worse
not all
weaker
not effect

stage & age
all stages
young only
late larvae and adults
adults only
old adults

comparison to other plates

comparison to known drugs

comparison to known mutants

- 27 -

Example 2**Profiling of a compound library (new compounds)**

5 To profile new compounds from a library, the general
profiling protocol is followed with the variations.
Compounds are profiled once in undiluted
concentration, the actual concentration being
dependent on the compound library in question but will
be between 0.01 mg and 1 mg of compound/10µl DMSO.

10 For compounds with a MW of 500 this calculates to 2-
200 mM stock. Dilution in 4ml agar would be at 5-500
µM. The high dose may create lots of unspecific effect
problems e.g. bacterial death and worm starvation.
15 Thus, if necessary the compounds are applied in a
second round at lower concentrations which are
dilutions in DMSO of 1/3, 1/10 and 1/30 of the
undiluted concentration. A concentration is finally
chosen for each compound which will allow a phenotype
20 profile to be established according to the standard
procedure.

Example 3**Profiling of known compounds (biotools, pharmacopoeia)**

25 To profile known compounds from a library the general
profiling protocol is followed with the following
variations. The stock solution is preferred as 100mM
in DMSO and the experiment is started *ab initio* with a
30 concentration series. The concentration series is used
as described below. In one series of concentrations 15
or so worms (for a reasonable number of short term
effects) are placed in the agar. In three series 1
worm each is placed on the agar to score a reasonable
35 number of progeny. Lost worms of the latter three
series of concentrations can be replaced from the

- 28 -

large pool where worms have been exposed to the compound in the same way. The following concentrations can be used:

5	conc.in 10 μ l drop	100mM	30mM	10mM	3 mM	1mM	0.3mM
	conc.in 4ml drop	100 μ M	300 μ M	100 μ M	30 μ M	10 μ M	3 μ M

Example 410 **Comparison of agar assay to drop assay**

A set of compounds from the pharmacopoeia have been profiled using the general protocol (all compounds were of known activity and are described in
15 Martindale: The Complete Drug Reference, 32nd edition, Pharmaceutical Press 1999). The plate drop assay was compared against standard of pouring compounds into the agar as described in literature which method is designated agar assay. In the drop assay as well as in
20 the agar assay, the compounds were added to the worm in a variety of concentrations, and the survival of the worm was scored as well as the phenotypic profile induced by the compound. The lowest concentration of a compound, still resulting in the death of the nematode
25 was designated minimal lethal dose. The maximal concentration of a compound that did not result in the death of the nematode was designated maximal nonlethal dose. The minimal concentration of a compound that still resulted in a measurable phenotype was
30 designated minimal effective dose. The concentrations of the compounds in the agar assay were compared to the concentrations in the drop assay. From this observation one may conclude that the newly described drop assay protocol turns out to be far more efficient
35 for most compounds. The following table lists the calculated concentration ratio needed to get the same

- 29 -

effect with the compound in the agar assay (in 2 ml agar) rather than the drop assay (in 4 ml agar).

Table 3:

5

10

15

20

Compound	Site	min. lethal dose	max. nonlethal dose	min. effective dose	average potency ratio
ketanserine	serotonin rec. agonist	>610			610
tamoxifen	estrogen rec. antagonist	204	304		254
fluoxetine	serotonin reuptake inh.	124	186		154
pancuronium	nicotinic antagonist			>100	100
methoxyphenylpiperazin	α -adrenorec. ligand	>48	>146	72	88
naloxone	opioid antagonist		>44	78	60
diheptylbipyridinium	ryanodine rec. antag.	20	30	36	28
W7	calmoduline antag.	20		10	14
thapsigargin	serca antagonist				14
physostigmine	cholinesterase inh.			8	8
lobeline	nicotinic rec. ligand			4	4
riluzole	glutamase release inh.	2	2	4	2
levamisole	acetylch. rec. antag			$\frac{1}{2}$	$\frac{1}{2}$
nicotine	acetylch. rec. antag			$\frac{1}{2}$	$\frac{1}{2}$

Minimal lethal dose: rate between the lowest concentration in which the compound is lethal to the worm in both assays
 Maximal non-lethal dose: rate between the highest concentration in which the compound is not lethal in both assays
 Minimal effective dose: rate between the lowest concentration in which the compounds results in a phenotype in both assays
 Average: average of the rates

- 30 -

Example 5**Preferred set of informative characteristics**

5 Worms exposed to a compound, carrying a mutation or
are transgenic are examined for the following 8
informative features/phenotypes:

1. Viability

10 Worms are examined for viability at all stages of the
life cycle, being embryogenesis, larval stages 1 to 4
and adulthood. Dead embryos are defined by not
hatching within 24h and dead worms are defined by not
moving, by lack of pharynx pumping, by sick or pale
15 appearance and by lack of response to mechanical
stimulation.

Method:

Embryonic lethality is measured by counting the amount
of unhatched worms after 24 hours (Elispot, Zeiss).
20 Counting of unhatched worms could also be automated
using the FANS device, described below. Viability of
larvae and adults is measured by dye uptake.

2. Life cycle

25 Progeny are examined for the length of the generation
cycle in comparison to control progeny (of a wild-type
worm). The stage of a synchronized progeny will be
compared to the stage of a synchronized control
progeny (N2, Bristol strain) after three days at 20°C.
30 The developmental stages can be distinguished by vulva
development, expression of stage-specific markers,
such as collagen IV, body length and transparency.

Method:

35 Measuring the body length of a population allows
determination of the actual stage in the life cycle

- 31 -

(For body shape measurement, see 3. Body shape).
Expression of stage-specific markers can be examined
using antibodies of the appropriate specificity, by
way of example an antibody that recognizes an antigen
5 on the surface of *C. elegans* L1 larvae has been
described by Hemmer et al., (1991) *J Cell Biol*,
115(5): 1237-47.

3. Body shape

10 Worm size is determined by measuring worm length and
worm diameter.

Method:

15 The body length of a synchronized progeny of adult
worms is compared to the body length of a synchronized
control progeny (N2, Bristol strain). Measurement of
body length can be achieved using a 'worm dispenser
apparatus' which is commercially available from Union
Biometrica, Inc, Somerville, MA, USA. This apparatus
20 has properties analogous to flow cytometers, such as
fluorescence activated cell scanning and sorting
devices (FACS). Accordingly, it may be commonly
referred to as a "FANS" apparatus, for fluorescence
activated nematode scanning and sorting device (FANS).
25 The FANS device enables the measurement of properties
of microscopic nematodes, such as size, optical
density, fluorescence, and luminescence.

30 Body size may also be measured via image analysis, in
which case the measurements recorded may include worm
diameter and deviation from the typical tube shape of
a wild-type worm.

4. Movement behaviour

35 The measurement of movement behaviour can include
measurement of the speed of movement, or of the

- 32 -

pattern of movement (e.g. direction) or both. A
wild-type worm moves in a sinusoidal way forward and
pauses or moves backward occasionally. Any deviation
from this wild-type pattern of movement can be scored
5 as a 'changed' characteristic.

Method:

An assay based on the following principles may be used
to determine the speed of movement of a worm culture:

10

Nematode worms that are placed in liquid culture will
move in such a way that they maintain a more or less
even (or homogeneous) distribution throughout the
culture. Nematode worms that are defective in
15 movement will precipitate to the bottom in liquid
culture. Due to this characteristic of nematode worms
as result of their movement phenotype, it is possible
to monitor and detect the difference between nematode
worms that move and nematodes that do not move.

20

Advanced multi-well plate readers are able to detect
sub-regions of the wells of multi-well plates. By
using these plate readers it is possible to take
measurements in selected areas of the surface of the
25 wells of the multi-well plates. If the area of
measurement is centralized, so that only the middle of
the well is measured, a difference in nematode
autofluorescence (fluorescence which occurs in the
absence of any external marker molecule) can be
30 observed in the wells containing nematodes that move
normally as compared to wells containing nematodes
that are defective for movement. For the wells
containing the nematodes that move normally, a low
level of autofluorescence will be observed, whilst a
35 high level of autofluorescence can be observed in the
wells that contain the nematodes that are defective in
movement.

- 33 -

In an adaptation of the movement assay, autofluorescence measurements can be taken in two areas of the surface of the well, one measurement in the centre of the well, and one measurement on the edge of the well. Comparing the two measurements gives analogous results as in the case if only the centre of the well is measured but the additional measurement of the edge of the well results in an extra control and somewhat more distinct results.

As an alternative to the above-described movement assay, specialist software such as SIMI Scout (designed for movement study of an athlete) may be used to determine speed of movement, deviation from sinusoidal movement and even the overall pattern of movement of the worm.

5. Mechanotransduction

Worms are examined for response to mechanical stimulation.

Method:

When the plate on which *C. elegans* are cultured is dropped wild-type worms react by enhanced movement and enhanced overall activity. The capability of a worm to respond to a mechanical stimulus is measured by the difference in speed of movement before and after stimulation.

6. Pharynx pumping

The phenotypes "Pumping frequency reduced, Pharynx pumping irregular" etc. describe the activity of the cyclic contraction of the pharynx muscles that occurs in a feeding adult about 3 times in a second. The contraction cycle can be described as the nearly simultaneously contraction of the corpus, anterior

- 34 -

isthmus, and terminal bulb, followed by relaxation.

Method:

5 The following pharynx pumping characteristics may be
analyzed by image analysis: The frequency of pumping
by counting the pharynx contraction. Pharynx
contraction can be measured visibly by the opening and
closing of the anterior corpus. The time of opened
anterior corpus and the diameter of the opened corpus
10 is used to measure hypercontraction, relaxation and
strength of a contraction.

The following is an example of a pumping assay which
allows measurement of the total efficiency of feeding
15 of a worm, which is related to pumping:

The pumping rate of the pharynx is measured indirectly
by adding a marker molecule precursor such as calcein-
AM to the medium and measuring the formation of marker
20 dye in the *C. elegans* gut. Calcein-AM is cleaved by
esterases present in the *C. elegans* gut to release
calcein, which is a fluorescent molecule. The pumping
rate of the pharynx will determine how much medium
will enter the gut of the worm, and hence how much
25 calcein-AM will enter the gut of the worm. Therefore
by measuring the accumulation of calcein in the
nematode gut, detectable by fluorescence, it is
possible to determine the pumping rate of the pharynx.

30 To perform the pharynx pumping screen with calcein-AM,
a concentration of between 1 and 100 μ M calcein-AM is
added into the medium. Preferably 5 to 10 μ M calcein-
AM is used. Fluorescence is measured using a multi-
well plate reader (Victor2, Wallac Oy, Finland) with
35 following settings: Ex/Em = 485/530.

- 35 -

7. Defecation

The defecation of *C. elegans* is a recurrent event comprising of the following steps: pBoc, aBoc and expulsion. Defecation in nematodes such as *C. elegans* is achieved by periodically activating a defined sequence of muscle contractions. These contractions are started in the anterior body wall muscles. At the zenith of the anterior body contractions the four anal muscles also contract. The four anal or enteric muscles are the two intestinal muscles, the anal depressor and the anal sphincter. In addition to this series of muscle contractions, specific neurons are also involved in the regulation of defecation, including the motor neurons, AVL and DVB.

Method:

In order to construct a phenotypic profile, well-fed adults are typically examined after one day for constipation. The time between two pBocs is also scored.

The rate of defecation of *C. elegans* can also be quantitatively measured using an assay based on the following principles:

The rate of defecation of nematodes such as *C. elegans* can be easily measured using a marker molecule which is sensitive to pH, for example the fluorescent marker BCECF. This marker molecule can be loaded into the *C. elegans* gut in the form of the precursor BCECF-AM which itself is not fluorescent. If BCECF-AM is added to nematode culture medium in the wells of a multi-well plate the worms will take up the compound which is then cleaved by the esterases present in the *C. elegans* gut to release BCECF. BCECF fluorescence is sensitive to pH and under the relatively low pH

- 36 -

conditions in the gut of *C. elegans* (pH<6) the compound exhibits no or very low fluorescence. As a result of the defecation process the BCECF is expelled into the medium which has a higher pH than the *C. elegans* gut and the BCECF is therefore fluorescent. The level of BCECF fluorescence in the medium (measured using a multi-well plate reader on settings Ex/Em=485/550) is therefore an indicator of the rate of defecation of the nematodes.

8. Fertility

A wild-type adult hermaphrodite *C. elegans* lays about 8 eggs per hour.

Method:

The amount of eggs laid by 20 hermaphrodite *C. elegans* during at least 60 min is counted. The amount of eggs may be counted by simple visual inspection or using a FANS device, described above.

Example 6

Comparison of profiles within a library

(daf-4 belongs to two pathways)


Mutant worms have been profiled according to the general profile protocol. Table 4 shows a summary of the profile, also called fingerprints, of one mutation of the indicated genes. Entries are binary with empty fields indicating a phenotype (deviation from negative control, here wild-type) not found assuming that it could have been measured. Any other entry including comments or quantitative data is read as measured phenotype in this binary scheme and indicated by *. The table lists only phenotypes that do have a

- 37 -

positive entry, not necessarily complete, leaving pages of empty fields alongside and arranged according to a particular enquiry. The upper half consists of the hierarchical categories "dauer formation phenotypes" and "body shape phenotypes" as well as their relevant sub-phenotypes. The lower part consists of a set of hierarchically unrelated phenotypes subsumed under the enquiry categories, "increased activity" and "decreased activity". The complete list of characteristics is to be found in Table 1.

The point of including the lower part is to show the principle of recording all observed phenotypes, that they can be used to distinguish similar phenotypic profiles in detail and that they can be arranged in order to make comparisons. In this case it is seen that the dichotomy of long versus short body length does not correlate to the dichotomy of increased versus decreased activity.

The upper part shows 5 genes (i.e. a mutation in that gene) affecting dauer formation as well as 5 genes affecting body shape in a particular combination. A mutation in one gene, daf-4, is unique in sharing the characteristics of both phenotypic groups. The following picture illustrates the phenotypic overlap as found by comparing entries in the phenotypic profiles.



35 From this overlap a hypothesis of a mechanistic link can be put forward for daf-4. In this particular case the mechanistic link is confirmed by the molecular

- 38 -

nature of the genes, which as far as known are all members of the TGF β pathway by sequence similarity:

		dbl-1 TGF β like ligand
5	daf-7 TGF β like ligand	sma-6 type I receptor
	daf-1 type I receptor	daf-4 type II receptor
	daf-4 type II receptor	sma-2 SMAD
	daf-3 SMAD	sma-3 SMAD
	daf-14 SMAD	sma-4 SMAD

10

The DAF-4 protein probably acts as a type II receptor in both pathways. The similarity of phenotypic profiles allows one to hypothesize mechanistic relationships in a manner analogous to sequence similarity of genes. For example a compound which induces the phenotypes: longer or shorter body length in combination with 2 or 3 of pale, thin and variable egg size, in worms exposed to it, is very likely to act on a protein of the TGF β pathway.

20

Table 4:

Phenotype	<i>daf-1</i>	<i>daf-7</i>	<i>daf-3</i>	<i>daf-14</i>	<i>daf-4</i> <i>e1364</i>	<i>sma-2</i> <i>e502</i>	<i>sma-3</i> <i>e491</i>	<i>sma-4</i> <i>e729</i>	<i>lon-1</i> <i>e185</i>	<i>lon-3</i> <i>e2175</i>
25 dauer formation	●	●	●	●	●					
constitutive dauer					
recovery defective					
body shape					●	●	●	●	●	●
30 short						
long									.	.
thin				
pale					

- 39 -

Phenotype	<i>daf-1</i>	<i>daf-7</i>	<i>daf-3</i>	<i>daf-14</i>	<i>daf-4</i> <i>e1364</i>	<i>sma-2</i> <i>e502</i>	<i>sma-3</i> <i>e491</i>	<i>sma-4</i> <i>e729</i>	<i>lon-1</i> <i>e185</i>	<i>lon-3</i> <i>e2175</i>
irregular egg size					•	•		•	•	•
increased activity					●		●	●	●	●
enhanced movement					•		•		•	
5 amplitude increased									•	
head movement enhanced							•	•	•	•
foraging behaviour increased					•			•		•
pharynx pumping enhanced							•		•	
constitutive pumping							•	•	•	
10 no egg retention									•	•
decreased activity						●				
lay still						•				
slow movement						•				
15 pharyngeal pumping reduced						•				

Example 7

20 **Comparison of phenotypes induced by acetylcholine
esterase inhibitors**

Wild type *C. elegans* adults have been exposed to
acetylcholine esterase inhibitors at various
25 concentrations. The worms have been profiled over two
generations, meaning four profiles have been
generated. All phenotypes from the phenotype list are
displayed that have been measured in this experiment.
Two phenotypes "loopy head movement" and "body dragged
30 by head" are shared by most of the esterase
inhibitors. This is called phenotype activity

- 40 -

relationship (PAR, by analogy to structure activity relationship SAR). The shared phenotypes are used to identify the action of a new compound. The unshared phenotypes are used to distinguish drugs or unravel side effects when these phenotypes are part of another PAR.

Table 5:

Phenotypes	Physostigmine	Neostigmine	Amibenonium	Tacrine	Galantamine	Trichlorfon
Thin	X					
Lay still	X					
Erratic	X					
Weak kinker		X				
Jerky				X		X
Enhanced head movement						X
Loopy head movement	X	X		X(L1)		X
Body dragged by head	X	X				X
Irregular touch response	X	X				
Reduced brood size	(X)					X
Delayed growth						X

Example 8

Comparison of phenotypes of mutations in the acetylcholine neurotransmission pathway

C. elegans adults and larval stages that are homozygous for the mutations *cha-1*, *unc-17*, *snt-1* and *cat-1* have been profiled, meaning fingerprints have been generated. All phenotypes from the phenotype list are displayed that have been scored in this

- 41 -

experiment. The phenotypes "small", "resistance to CHA inhibitors (Ric)", "slow pumping" and "slow growth" are shared. This is called phenotype activity relationship (PAR, in analogy to structure activity relationship SAR). The shared phenotypes are used to identify genes in a pathway. The unshared phenotypes are used to distinguish these genes or unravel further functions in parallel or new pathways when these phenotypes are part of another PAR. The fingerprint of *cat-1* is different because this gene is involved in the dopamine pathway.

Table 6:

Phenotype	<u><i>cha-1</i></u> ChAT (synthesis)	<u><i>unc-17</i></u> VchAT (ACh- transporter)	<u><i>snt-1=ric-2</i></u> Synaptotag min homolog	<u><i>cat-1</i></u> VMAT (monamine- transporter)
Coiler	X	X		
Small	X	X	X	
Slow growth	X	X	X	
Ric	X	X	X	
Slow pumping	X	X	X	
Jerky when backing	X			X
Low ChAT level	X			
Poor male turning				
Enhanced foraging behaviour				
Enhanced foraging behaviour				
Defecation defects				
Shrinker-uncs				

- 42 -

Example 9

Method to profile an intervention (mutation, compound etc)

- 5 Profiling a mutation in the gene *unc-17* that affects transportation of acetylcholine.

10 In the literature this phenotype is described, concerning movement, body size and feeding, as severe coiler, being rather small and thin and has only slow, irregular pumping of the pharynx (Riddle et al., "C. elegans II" Cold Spring Harbor Laboratory Press, 1997). By systematically describing *unc-17* the resulting fingerprint unravels more details and new

15 properties: Concerning movement, body size and feeding the phenotypes strong coiler, spiralling inwards posteriorly, curly jerky and moves better forward, being small have been profiled. In addition defects in the sensory system, defecation and reproductive system

20 have been found, in detail: the touch response is gone, constipation, aberrant defecation cycle (aBoc) and egg laying defective (no egg retention).

25 **Example 10**

Method to add biological information to a particular phenotype

30 One phenotype of the mutation *unc-4* is "coiler" (looks like a snail). The fingerprint of *unc-4* adds for "coiler" the details "ventral side out" and "spiralling inwards posteriorly". This occurs when a set of neurons that control the forward movement of the ventral part of the worm (VA2 - VA10) gets the

35 same input than another set of neurons that controls the backward movement of the ventral part (VB2 -

- 43 -

VB10).

In this case the ventral muscles get contradicting signals and only the dorsal muscles contract properly. The result is a coiler that has only the ventral side outwards. We explain most of the phenotypes as consequence of a mislead process, here synaptic input.

10 **Example 11**

Comparison of phenotypes induced by compounds acting on GABAergic neurotransmission

15 Wild-type *C. elegans* adults have been exposed to GABA agonists (Muscimol) and GABA antagonists (Ivermectin and Fipronil) at various concentrations. Worms have been profiled and the scored phenotypes are displayed as fingerprints.

20 In addition, two mutations in the GABAergic pathway have been profiled and compared with the compound induced phenotypes: *unc-25* encodes for the decarboxylase and *unc-49* encodes for a GABA receptor.

25 The phenotype "shrinker" is present in all fingerprints (see Table dark grey). This phenotype is used as marker or diagnostic phenotype to identify activity of a compound or gene in the GABAergic pathway. There are further phenotypes only shared by
30 some compounds and mutants (see Table light grey). These phenotypes are used to build a phenotype activity relationship (PAR).

35 The shared phenotypes are used to identify the action of a new compound when "shrinker" cannot be used or to reveal more details on a compound action. For example,

- 44 -

all compounds and *unc-25* fingerprints contain constipation phenotypes but not the fingerprint of *unc-49*, although GABA is used for the defecation process. This is coincident with earlier findings that the UNC-49 gene product is not required for defecation.

These results may indicate the existence of another yet unknown GABA receptor in *C. elegans*. The unshared phenotypes are used to unravel toxic side effects or other mode of actions.

Table 7:

Phenotypes	Muscimol	Ivermectin	Fipronil	<i>unc-25</i>	<i>unc-49</i>
Pale	x	x		X	
Motionless (paralyzed) I	x	x			
Nearly motionless	x	x			
No movement but motion II	x		x	X	x
Little movement	x		x	X	x
Slow movement III	x		x	X	
Enhanced movement V					
Stiff rods					
Loose rods	x	x			
Rigid paralysis (hypercontracted)					
Flaccid paralysis (relaxed)	x	x			
Bent body, jerky body, abnormal Omega appearance			x	(x)	
Enhanced foraging			x		x
Shrinker before movement	x		x	X	
Shrinker	x	x		X	x
No pumping	x	x			
Weak pumping					
Pumping frequency reduced		x	x		
Pumping frequency enhanced					
Pumping irregular	x				
Constipation		x	x	X	
Foregut filled/enlarged			x		
Hindgut weak constipated		x	x	X	
Hindgut strong constipated				X	
Defecation cycle defective	x	x	x	x	
(time: pBoc)					
Weak expulsion				x	
No expulsion				x	
No egg retention (12-cell stage)					
Weak egg laying defect (comma)					
Strong egg laying defect (pretzel)		x	x		
Blatant worms			x		
Bags of worms			x		

- 45 -

Example 12**Definition of body shape phenotypes**

Aberrations of the body shape of *C. elegans* can be the
 5 result of mutations in a vast amount of genes. These
 genes may be required directly for the formation of
 the hypodermis, the hydroskeleton and the correct
 patterning of the worm body plan, e.g., collagen or
 even-skipped. They could be involved in the control of
 10 growth or metabolism like genes of the TGF β pathway
 or genes required for feeding. Eventually, mutations
 in certain genes that cause primary defects, e.g.,
 absence of head muscle, cause secondary defects in the
 body shape like dystrophy in the head region.
 15 Body shape phenotypes are all visible or measurable
 deviations of the body shape, colour and content.
 Phenotypes are comparatively measured against wild-
 type (N2, Bristol strain) and scored as deviation of
 wild type in the corresponding developmental stage,
 20 sex and preparation. The scored phenotype comes with
 the percentage of worms positive for that phenotype
 within a population.

Table 8: Scientific definition of body shape
 25 phenotypes. The phenotypes listed in the left column
 are described and defined in the right column. Some
 phenotypes are derived from the classical worm jargon
 like "dumpy", which is still shorter than "short and
 thick worm".

30

PHENOTYPE	DEFINITION
Proportion abnormal	
Short	Body length less than wild type.
Long	Body length more than wild type.
Thin	Body diameter less than wild type.
Thick	Body diameter more than wild type.

35

- 46 -

Dumpy	Body length less but body diameter more than wild type.
Spindle-shaped	body diameter is more for only a restricted region of the body.

Head defects

Hypertrophy of the head	Regions of the head are thickened. This additional tissue is part of the head and enclosed by the hypodermis.
Extensions of head	Small hypertrophied regions of the head.
Notched head	Extensions, protrusions on the dorsal side of the head.
Hammer head	Extensions at the head tip resemble a hammer like appearance.
Dystrophy of the head	Regions of the head are thinned due to missing tissue.
Swollen	The head looks like a balloon.
Rounded	The tip of the head is rounded.
Tapering	The tip of the head is tapering.
Vacuoles only in head	Vacuoles visible in the head but not in the rest of the body.
Only head bent	The head is held most of the time in a bent position. In extreme cases the worm looks like a walking stick.
Autodecapitation	The head/body connection is thinner, which results occasionally in an autodecapitation due to a body wall muscle contraction.

Body defects

Scrawny	Worm is shorter, thinner, pale and sick.
Hypertrophy of body	Regions of the body are thickened. This additional tissue is part of the body and is enclosed by the hypodermis.
Extensions	Small hypertrophied regions of the body.
Humpback	Extensions, protrusions on the dorsal side of the body. The counterpart, extensions on the ventral side of the body, would be scored as "multi vulva" in the section "Vulva". The distinction between a non vulva-like extension versus a vulva-like extension will be made with a high power microscope.
Truncated body	Part of the body is missing.
Withered body	Part of the body is thinned.
Twisted	Twisted body. The rotation along the anterior-posterior body axis can be seen by the twisted gut/gonad tube or because the vulva and the rectum are not orientated in the same (ventral) direction.
Fat	Worm is thicker and darker than wild type.
Pale	Worm is brighter than wild type.
Pale with dark spots	Worm is brighter than wild type and contains dark spots.
Clear	Worm is nearly transparent.
Full of vacuoles	Worm contains more vacuoles than wild type. Vacuoles have a darker or opal appearance and resemble little moon craters.
Fluid-filled	Liquid flows all over the body.

- 47 -

Poured out	Contents of the worm like the gonad is released through th vulva.
Burst	Dead worm with bursted body shape.

Tail defects

Only tail truncated	Blunt body end; whipe is missing.
Tail shape aberrant	Tail or tail whipe is kinked, shortened or thickened.
Knob-like	Tail whipe has knob-like structures.

Cuticle defects

Blistered	Fluid-filled transparent blisters separated by the hypodermis outside on the body. Clearly different from extensions.
Molting defective	More worms are caught in their old skin like the sloughing of a snake.

10

It is possible to score body shape phenotypes by image acquisition followed by image analysis. The advantage in the automation of the profiling procedure is the quantification of the strength of a phenotype or the presence of the phenotype in a population. A disadvantage is that the procedure for analysing an image for every possible phenotype may be more elaborate than simply scoring by eye. Furthermore, certain details are difficult to access by video analysis e.g., blister versus protrusions.

Table 10: list of scientific body shape phenotypes, together with their corresponding technical definitions , in terms of characteristics which can be comparatively measured relative to wild-type characteristics using automated measuring apparatus.

Scientific phenotype	Technical definition	Technical phenotype
Proportion abnormal		
Short	Body length less than wild type	Short
Long	Body length more than wild type	Long
Thin	Body diameter less than wild typ	Thin

30

- 48 -

Thick	Body diameter more than wild type	Thick
Dumpy		<i>Disappears</i>
Spindle-shaped		<i>Disappears</i>

Head defects

5	Hypertrophied head	Total head volume has increased	Hypertrophied head
	Extensions on head	Head will be subdivided in n trapezes (or n slices). The diameter of different trapezes can be compared pairwise. The deviation of the diameter can also be located to one side	Extensions on head
	Notched head		Extensions only on one side
	Hammer head		Extensions are pairwise
	Dystrophied head	Total head volume has decreased	Dystrophied head
10	Swollen		<i>Disappears</i>
	Rounded	In the tip trapeze the top diameter is increased	Rounded
	Tapering	The diameter of the tip trapezes are decreased	Tapering
	Vacuoles only in head		<i>Disappears</i>
	Only head bent	The head is most of the time in a certain position that can be measured by an average angle between tip and head/body connection	Tip of head is more often in one position
15	Autodecapitation		<i>Disappears</i>

Example 13**Use of GFP in profiling *C. elegans***

20

A lot of features of *C. elegans* as described in Table 1 can be easily monitored, either automatically by image analysis, microtiter plate readers, or visual means, e.g. by normal microscopy or by Nomarski

25 microscopy. Some features of *C. elegans* are more difficult to visualize. For these characteristics transgenic animals expressing a marker gene are very useful. Moreover, even for characteristics that are rather easily to score, the use of a nematode

30 expressing a marker gene, such as GFP, LacZ, or luciferase, enhances the fingerprinting of *C. elegans*.

- 49 -

The *C. elegans* can be a wild type, a mutant, or a strain subjected to a compound or environmental stress, or a combination of those.

5 *C. elegans* mutant *unc-23* has a fingerprint, which comprises "jerky movement", "tend to coil", "bent head" and "egl". Expressing GFP in the muscle cells of the animal could result in identification and scoring of additional characteristics such as "improperly
10 folded muscles", and/or "detached muscles in head region", and/or "no muscles in head region", and/or "defective muscle attachment", and/or "vulva muscle defects" (data not shown).

15 Similarly, *C. elegans* mutant *unc-71* has a fingerprint which comprise "reduced movement", "weak amplitude", "strong kinker", and "slightly egl". When introducing GFP in the neurons of the animals no apparent extra fingerprint features were observed. A closer look at
20 the neurons of this mutant worm revealed at least following extra phenotypes: "fasculation defects", "VD/DC connection defects" (data not shown).

GFP-phenotypes are hence very important in allowing
25 phenotypes which are not otherwise visible to be measurable with Nomarski or dissection microscopy. GFP-phenotypes are further important in the pinpointing of defects to certain tissues and cells, and moreover GFP-phenotypes are important in
30 distinguishing between similar defects with different causes.

- 50 -

Claims:

1. A method of constructing a library of phenotypic profiles of nematode worms which comprises
5 the steps of:

(a) providing a worm having a defect in at least one gene,

10 (b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,

(c) systematically scoring a plurality of any
15 said changed characteristics to establish a characteristic phenotypic profile associated with said defect,

(d) simultaneously or sequentially repeating
20 steps (a) to (c) in respect of each of a plurality of worms each of which has a different defect, and

(e) collating the phenotypic profiles so obtained
25 into a library of said profiles.

2. A method as claimed in claim 1 wherein in step (c) at least three changed characteristics are scored.

30 3. A method as claimed in claim 1 or claim 2 wherein in step (c) at least six changed characteristics are scored.

35 4. A method as claimed in any preceding claim wherein in step (c) at least ten characteristics are scored.

- 51 -

5. A method as claimed in any preceding claim wherein said worm is *Caenorhabditis elegans*.

6. A method as claimed in any preceding claim
5 wherein steps (a) to (c) are carried out in respect of substantially every gene in the worm genome.

7. A method as claimed in any preceding claim
10 which includes the step of manipulating said worm to generate said defect in said at least one gene.

8. A method as claimed in any preceding claim wherein said defect is selected from the absence of expression of said gene, the reduction in expression
15 of said gene, the over-expression of said gene, the expression of a functionally defective protein, the expression of a truncated protein, the misexpression of a protein, the ectopic misexpression of a protein, the expression of a protein of altered stability or
20 the alteration of gene expression as a function of time.

9. A method as claimed in claim 7 or 8 wherein said manipulation is carried out on wild-type *C. elegans* or a selected mutant thereof.
25

10. A method as claimed in claim 9 wherein said selected mutant harbours multiple mutations.

30 11. A method as claimed in claim 7 or 8 wherein said manipulation is carried out on *C. elegans* carrying a reporter gene.

12. A method as claimed in claim 11 wherein said
35 reporter gene is LacZ or green fluorescent protein (GFP).

- 52 -

13. A method as claimed in any one of claims 7 to 12 wherein said manipulation is carried out on a transgenic *C. elegans*.

5 14. A method as claimed in claim 13 wherein said transgenic *C. elegans* expresses a human gene.

15. A method as claimed in claim 14 wherein said human gene is a known drug target.

10 16. A method as claimed in claim 14 or claim 15 wherein said human gene is one associated with a human disease.

15 17. A method as claimed in claim 14 or 15 wherein said human gene is a candidate human disease gene.

20 18. A method as claimed in any of claims 7 to 17 wherein said manipulation is carried out on only a sub-set of *C. elegans* cells.

25 19. A method as claimed in any preceding claim wherein changed characteristics in said worm carrying said defect compared to a worm that does not carry said defect are identified by light microscopy, differential interference contrast optics, fluorescence microscopy, immunochemical detection or spectrophotometric detection, radiation detection, calorimetric detection, fluorescence detection or luminescence detection.

30 20. A method as claimed in any preceding claim wherein changed characteristics in said worm carrying said defect compared to a worm that does not carry said defect are identified by a pH change or a change in electrical potential.

- 53 -

21. A method as claimed in any preceding claim wherein said plurality of changed characteristics are scored in a predetermined order to generate said phenotypic profile.

5

22. A method as claimed in any preceding claim wherein the scoring of said plurality of changed characteristics is repeated at predetermined intervals of time.

10

23. A method as claimed in any preceding claim wherein said phenotypic profiles are stored electronically.

15

24. A method as claimed in any preceding claim wherein at least one of said plurality of characteristics is selected from the list shown in Table 1.

20

25. A method as claimed in any one of the preceding claims wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

25

26. A method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

30

(a) exposing a worm to a compound,

(b) measuring any changes in identifiable characteristics of said worm as a result of exposure to said compound,

35

(c) systematically scoring a plurality of any

- 54 -

said changed characteristics to establish a phenotypic profile associated with said compound,

5 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different compounds and

(e) collating the phenotypic profiles so obtained into a library of said profiles.
10

27. A method as claimed in claim 26 wherein in step (c) at least three changed characteristics are scored.

15 28. A method as claimed in claim 27 wherein in step (c) at last six changed characteristics are scored.

20 29. A method as claimed in claim 28 wherein in step(c) at least ten changed characteristics are scored.

25 30. A method as claimed in any one of claims 26 to 29 wherein said nematode worm is *C. elegans*.

31. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds has a known pharmacological activity.

30 32. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds is one which is known to interact with a particular biochemical pathway.

35 33. A method as claimed in any one of claims 26 to 30 wherein each of said plurality of different compounds has no known pharmacological activity or

- 55 -

biochemical interaction.

34. A method as claimed in any one of claims 26
to 30 wherein each of said plurality of different
5 compounds is from a combinatorial library.

35. A method as claimed in any one of claims 26
to 34 wherein said worm to which said compound is
exposed is wild-type *C. elegans* or a selected mutant
10 thereof.

36. A method as claimed in claim 35 wherein said
selected mutant harbours multiple mutations.

15 37. A method as claimed in any one of claims 26
to 34 wherein said worm to which said compound is
exposed is *C. elegans* carrying a reporter gene.

38. A method as claimed in claim 37 wherein said
20 reporter gene is LacZ or GFP.

39. A method as claimed in any one of claims 26
to 38 wherein said worm to which said compound is
exposed is a transgenic *C. elegans*.

25 40. A method as claimed in claim 39 wherein said
transgenic *C. elegans* expresses a human gene.

41. A method as claimed in claim 40 wherein said
30 human gene is a known drug target.

42. A method as claimed in claim 40 wherein said
human gene is one associated with a human disease.

35 43. A method as claimed in claim 40 wherein said
human gene is a candidate disease gene.

- 56 -

44. A method as claimed in any one of claims 30 to 43 wherein said worm is exposed to said compound by feeding the worm on bacteria which have been exposed to said compound.

5

45. A method as claimed in claim 44 wherein said bacteria are *E. coli*.

46. A method as claimed in any one of claims 26 to 45 wherein said compound is linked to another compound or carrier substance.

10

47. A method as claimed in anyone of claims 26 to 46 wherein any changed characteristics in said worm resulting from exposure to said compound are identified by light microscopy, differential interference contrast optics, fluorescence microscopy, immunochemical detection, spectrophotometric detection, radiation detection, colorimetric detection, fluorescence detection or luminescence detection.

15

20

48. A method as claimed in any one of claims 26 to 47 wherein any changed characteristics in said worm resulting from said compound are identified by a pH change or a change in electrical potential.

25

49. A method as claimed in any one of claims 26 to 48 wherein said plurality of changed characteristics are scored in a predetermined order to generate said profile.

30

50. A method as claimed in any one of claims 26 to 49 wherein the scoring said plurality of changed characteristics is repeated at predetermined time intervals.

35

- 57 -

51. A method as claimed in any one of claims 26 to 50 wherein said scoring of changed characteristics is carried out using essentially the same scoring protocol as used in a method in accordance with any one of claims 1 to 25.

52. A method as claimed in any one of claims 26 to 51 which comprises the further step of storing the said phenotypic profiles electronically.

53. A method as claimed in any one of claims 26 to 52 wherein at least one of said plurality of characteristics is selected from the list shown in Table 1.

54. A method as claimed in any one of claims 26 to 53 wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

55. A method of constructing a library of phenotypic profiles of nematode worms which comprises the steps of:

(a) exposing a worm to an environmental change,

(b) measuring any changes in identifiable characteristics as a result of said environmental change,

(c) systematically scoring a plurality of any said changed characteristics to establish a Characteristic phenotypic profile associated with said change,

- 58 -

(d) simultaneously or sequentially repeating steps (a) to (c) for each of a plurality of different environmental changes and (e) collating the phenotypic profiles so obtained into a library of said profiles.

5

56. A method as claimed in claim 55 wherein in step (c) at least three changed characteristics are scored.

10

57. A method as claimed in claim 56 wherein in step (c) at least six changed characteristics are scored.

15

58. A method as claimed in claim 57 wherein in step (c) at least ten changed characteristics are scored.

20

59. A method as claimed in any of claims 55 to 58 wherein said environmental change is a change in the pH to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a different pH.

25

60. A method as claimed in any one of claims 55 to 58 wherein said environmental change is a change in the osmolarity to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a different osmolarity.

30

61. A method as claimed in any one of claims 55 to 58 wherein said environmental change is a change in the temperature to which the worm is exposed and in step (d) each of the plurality of environmental changes comprises a change in temperature.

35

62. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises

- 59 -

exposure to radiation and in step (d) each of said plurality of environmental changes comprises a different level of radiation.

5 63. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises exposure to a virus and in step (d) each of said plurality of environmental changes comprises exposure to a different virus.

10

 64. A method as claimed in any one of claims 55 to 58 wherein said environmental change comprises exposure to a bacterium and in step (d) each of said plurality of environmental changes comprises exposure to a different bacterium.

15

 65. A method as claimed in any one of claims 55 to 64 wherein said worm is *C. elegans*.

20 66. A method as claimed in any one of claims 55 to 65 including a further feature as defined in any one of claims 5 to 54.

25 67. A method as claimed in any one of claims 55 to 66 wherein said scoring of changed characteristics is carried out using essentially the same scoring protocol as used in a method in accordance with claims 1 to 54.

30 68. A method as claimed in any one of claims 55 to 67 wherein step (b) comprises measuring changes in two or more characteristics selected from the group consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

35

 69. A method of constructing a multiple library

- 60 -

of phenotypic profiles of nematode worms which method comprises carrying out all of the methods of claims 1, 26 and 55.

5 70. A method as claimed in claim 69 wherein step (b) of the method of at least one of claims 1, 26 and 55 comprises measuring changes in two or more characteristics selected from the group consisting of:
10 viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

 71. A method of determining the mode of action of a compound which method comprises the steps of;
15

(a) exposing a nematode worm to said compound

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure
20 to said compound,

(c) systematically scoring a plurality of changed characteristics to establish a phenotypic profile associated with said compound and
25

(d) comparing said phenotypic profile with a library of reference phenotypic profiles wherein said library of reference profiles is obtainable by carrying a method in accordance with any of claims 1
30 to 70.

 72. A method of determining whether a compound or combination of compounds interacts with a particular gene or biochemical pathway which method
35 comprises the steps of;

(a) exposing a nematode worm to said compound or

- 61 -

combination of compounds

5 (b) measuring any changes in identifiable characteristics of said worm as a result of said exposure,

10 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile associated with said compound or combination of compounds, and

15 (d) comparing said profile with a library of reference profiles said library of reference profiles being obtainable by carrying out the method of any one of claims 1 to 70.

20 73. A method of finding an alternative treatment for a human disease which method comprises the steps of:

(a) exposing a nematode worm to a candidate compound,

25 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

30 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

35 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 31.

74. A method of finding a biochemical pathway in

- 62 -

which a compound known to have pharmacological activity acts which method comprises the steps of:

5 (a) exposing a nematode worm to the known compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

10

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound, and

15 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 32.

20 75. A method of finding a potential new medicinal indication for a compound of known pharmaceutical activity which method comprises the steps of:

25 (a) exposing a nematode worm to the known compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

30

(c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

35

(d) comparing said profile with a library of reference profiles, said library of reference profiles

- 63 -

being obtainable by carrying out a method in accordance with any one of claims 1 to 70.

5 76. A method as claimed in claim 75 wherein said library of reference profiles is obtainable by carrying out a method in accordance with any one of claims 24 to 26.

10 77. A method of identifying the mechanism of action of any side effects associated with a compound of known pharmaceutical activity which method comprises the steps of;

15 (a) exposing a nematode worm to the known compound,

20 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

25 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound and

30 (d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by carrying out a method in accordance with claim 32 and/or any of claims 1 to 25.

35 78. A method of attributing a particular gene to a particular biochemical pathway in *C. elegans* which method comprises the steps of:

 (a) exposing a nematode worm to a compound known to operate in a particular biochemical pathway,

 (b) measuring any changes in the identifiable

- 64 -

characteristics of said worm as a result of exposure to said compound

5 (c) systematically scoring a plurality of any changed characteristics to establish a phenotypic profile for said compound, and

(d) comparing said, profile with a library of reference phenotypic profiles said library of
10 reference profiles being obtainable by carrying out a method in accordance with any one of claims 1 to 25.

79. A method as claimed in any of claims 71 to 78 wherein said nematode worm is selected from wild-
15 type *C. elegans*, a mutant *C. elegans* comprising one or more mutations, a *C. elegans* carrying a reporter gene or a transgenic *C. elegans*.

80. A method as claimed in claim 79 wherein said
20 transgenic *C. elegans* expresses a human gene.

81. A method as claimed in any one of claims 71 to 80 wherein step (b) comprises measuring changes in two or more characteristics selected from the group
25 consisting of: viability, life cycle, body shape, movement behaviour, mechanotransduction, pharynx pumping, defecation and fertility.

82. A method for elucidating biochemical
30 pathways in a nematode worm which method comprises the steps of:

(a) generating a defect in at least one gene in said worm,
35

(b) measuring any changes in identifiable characteristics of said worm compared to a worm

- 65 -

without said defect,

(c) systematically scoring a plurality of any
said changed characteristics to establish a phenotypic
5 profile for said defect, and

(d) comparing said profile with a library of
reference phenotypic profiles, said library of
references profiles being obtainable by carrying out a
10 method in accordance with any one of claims 1 to 25.

83. A method as claimed in claim 82 wherein said
nematode worm is selected from wild-type *C. elegans*, a
mutant *C. elegans* comprising one or more mutations, a
15 *C. elegans* carrying a reporter gene or a transgenic *C. elegans*.

84. A method as claimed in claim 82 wherein said
defect is selected from the absence of expression of
20 said gene, the reduction in expression of said gene,
the expression of a functionally defective protein,
the expression of a truncated protein, the
misexpression of a protein, the ectopic misexpression
of a protein, the expression of a protein of altered
25 stability or the alteration of gene expression as a
function of time.

85. A method as claimed in any one of claims 82
to 84 wherein at least three, preferably at least six
30 and more preferably at least ten changed
characteristics are scored.

86. A method as claimed in any of claims 82 to
85 which includes the features described in any one of
35 claims 19 to 25.

87. A method of constructing a library of

- 66 -

nematode worms which method comprises the steps of:

(a) providing a worm having a defect in at least one gene.

5

(b) measuring any changes in identifiable characteristics of said worm compared to a worm without said defect,

10

(c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said defect,

15

(d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of worms, and

20

(e) producing a library of said worms each identifiable by their phenotypic profiles.

88. A method as claimed in claim 87 wherein said phenotypic profiles are collated into a library.

25

89. A method as claimed in claim 87 and 88 comprising any one of the features described in any one of claims 2 to 25.

30

90. A method of constructing a library of nematode worms which method comprises the steps of:

(a) exposing a worm to a compound,

35

(b) measuring any changes in identifiable characteristics of said worm as a result of exposure to said compound,

- 67 -

(c) systemically scoring a plurality of any said changed characteristics to establish a phenotypic profile associated with said compound,

5 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different compounds, and producing a library of said worms each identifiable by their phenotypic profiles.

10 91. A method as claimed in claim 90 wherein said phenotypic profiles are collated into a library.

92. A method as claimed in claim 90 or 91 comprising any one of the features disclosed in any
15 one of claims 27 to 54.

93. A method of constructing a library of nematode worms which method comprises the steps of:

20 (a) exposing a worm to an environmental change,

(b) measuring any changes in identifiable characteristics as a result of said environmental change,

25 (c) systematically scoring a plurality of any said changed characteristics to establish a characteristic phenotypic profile associated with said change,

30 (d) simultaneously or sequentially repeating steps (a) to (c) in respect of each of a plurality of different environmental changes, and

35 (e) producing a library of said worms each identifiable by their phenotypic profile.

94. A method as claimed in claim 93 wherein said

- 68 -

phenotypic profiles are collated into a library.

95. A method as claimed in claim 93 or claim 94 comprising any one of the features disclosed in any
5 one of claims 56 to 70.

96. A method of determining the mode of action of a compound which method comprises the step of:

10 (a) exposing a nematode worm to said compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,
15

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile associated with said compounds, and

20 (d) comparing said phenotypic profile with the library of phenotypic profiles obtainable by the method of any one of claims 88, 91 or 94.

97. A method of determining whether a compound
25 or a combination of compounds interacts with a particular gene or biochemical pathway which method comprises the steps of:

(a) exposing an nematode worm to said compound or
30 combination of compounds,

(b) measuring any changes in identifiable characteristics of said worm as a result of said exposure,
35

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic

- 69 -

profile associated with said compounds or combination of compounds, and

- 5 (d) comparing said phenotypic profile with a library of reference profiles wherein said library of reference profiles is obtainable by the method of any one of claims 88, 91 or 94.

10 98. A method of finding an alternative treatment for a human disease which method comprises the steps of:

(a) exposing an nematode worm to a candidate compound,

15 (b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

20 (c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

25 (d) comparing said profile with a library of referenced profiles, wherein said library of referenced profiles is obtainable by carrying out the method in accordance with any one of claims 88, 91 or 94.

30 99. A method of finding a biochemical pathway in which a compound known to have pharmacological activity acts which method comprises the steps of:

35 (a) exposing a nematode worm to the known compound, measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

- 70 -

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

5 (c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

(d) comparing said profile with a library of
10 reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

100. A method of finding a potential new
15 medicinal indication for a compound of known pharmaceutical activity which method comprises the steps of:

(a) exposing an nematode worm to the known
20 compound,

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

25

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

(d) comparing said profile with a library of
30 reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

35 101. A method of identifying the mechanism of action of any side effects associated with a compound of known pharmaceutical activity which method

- 71 -

comprises the steps of:

(a) exposing a nematode worm to the known compound,

5

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

10

(c) systematically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

15

(d) comparing said profile with a library of reference profiles, said library of reference profiles being obtainable by the method of any one of claims 88, 91 or 94.

20

102. A method of attributing a particular gene to a particular biochemical pathway in *C. elegans* which method comprises the steps of:

(a) exposing a nematode worm to a compound known to operate in a particular biochemical pathway,

25

(b) measuring any changes in the identifiable characteristics of said worm as a result of exposure to said compound,

30

(c) systemically scoring a plurality of any said changed characteristics to establish a phenotypic profile for said compound, and

35

(d) comparing said profile with a library of reference phenotypic profiles, said library of reference profiles being obtainable by carrying out the method in accordance with any one of claims 88, 91

- 72 -

or 94.

103. A method as claimed in any one of claims 96
to 102 wherein said nematode worm is selected from
5 wild-type *C. elegans*, a mutant *C. elegans* comprising
one or more mutations, a *C. elegans* carrying a
reporter gene or a transgenic *C. elegans*.

104. A method as claimed in claim 103 wherein
10 said transgenic *C. elegans* expresses a human gene.

105. A method of establishing a phenotypic
profile for a nematode worm which method comprises
measuring and scoring at least three, preferably at
15 least six and more preferably at least ten
characteristics of said worm which are not exhibited
by wild-type worms.

106. A method as claimed in claim 105 wherein
20 said characteristics not exhibited by wild-type worms
are selected from the list shown in Table 1.

107. A method as claimed in claim 105 or claim
106 which comprises measuring and scoring changes in
25 two or more characteristics selected from the group
consisting of: viability, life cycle, body shape,
movement behaviour, mechanotransduction, pharynx
pumping, defecation and fertility.

108. A method as claimed in any one of claims 105
30 to 107 wherein said phenotypic profile is established
for a nematode worm which is selected from a worm
having one or more mutations, a worm which has been
exposed to a compound or combination of compounds, a
35 transgenic worm, a worm carrying a reporter gene or a
worm which has been exposed to an environmental
change.

- 73 -

109. A method as claimed in claim 108 wherein
said transgenic worm comprises a human gene.

5 110. A method as claimed in claim 108 wherein
said compound has known pharmacological activity.

10 111. A method as claimed in claim 108 wherein
said compound is known to be active in a particular
biochemical pathway.

112. A method as claimed in claim 108 wherein
said compound or combination of compounds is from a
combinatorial library of compounds.

15 113. A compound which has potential therapeutic
activity in a mammal which has been identified in a
method as claimed in any one of claims 71 to 81 or 96
to 104.

20 114. A library of nematode worms obtainable by a
method as claimed in any one of claims 87 to 95.

25 115. A library as claimed in claim 114 wherein
said nematode worm is *C. elegans*.


FIG. 1.

FIG. 2.

